NEURODIVERGENCE, INTERSECTIONALITY, AND EATING DISORDERS: A LIVED EXPERIENCE-LED NARRATIVE REVIEW [PREPRINT]

This manuscript has been submitted for publication

Neurodivergence, intersectionality, and eating disorders: A lived experience-led narrative review

Laurence Cobbaert ^{1, 2*}, Anna Rose ^{3, 2}, Rosiel Elwyn ⁴, Scout Silverstein ⁵, Kai Schweizer ⁶, Elysia Thomas ^{7, 2}, Jane Miskovic-Wheatley ⁸

*Corresponding author: 1.cobbaert@unsw.edu.au

- ¹ University of New South Wales, Australia
- ² Eating Disorders Neurodiversity Australia, Australia
- ³ Bond University, Australia
- ⁴ Thompson Institute, University of the Sunshine Coast, Australia
- ⁵ Project HEAL, New York, United States of America
- ⁶ Telethon Kids Institute, University of Western Australia, Australia
- ⁷ Independent researcher, Australia
- 8 InsideOut Institute, University of Sydney, Australia

ABSTRACT

Autistic people and those with attention deficit hyperactivity disorder are at a high risk of developing an eating disorder. While there is limited evidence on the relationship between other forms of neurodivergence and eating disorders, research suggests associations between giftedness, intellectual disability, Tourette's syndrome, and disordered eating. Factors underlying disordered eating and/or eating disorder risk for neurodivergent people are multifaceted and complex, encompassing a wide range of intertwined psychosocial, environmental, and biological processes. Moreover, research shows that neurodivergent individuals experience poorer treatment outcomes compared to neurotypical individuals. However, there is a paucity of research in this area overall. More specifically, lived experience-led research remains rare, despite its critical role for improving individualised eating disorder care, as well as mental healthcare more broadly. Indeed, the importance of eating disorder care individuation is increasingly being recognised, particularly within the context of neurodivergence, given the heterogeneous experiences and support needs of neurodivergent people affected by disordered eating and/or eating disorders. Furthermore, despite documented overlaps between various forms of neurodivergence (e.g., co-occurring autism and attention deficit hyperactivity disorder), research looking at eating disorders in the context of neurodivergence through a transdiagnostic perspective is scarce. This lived experience-led narrative review aims to shed light on the intersectional factors underlying high disordered eating and/or eating disorder risk for neurodivergent individuals. First, an overview of prevalence data is provided, followed by a thematic framework identifying factors underlying disordered eating and/or eating disorder risk in relation to neurodivergence. A critical appraisal of current eating disorder research and care is then offered before suggestions for neurodiversity-affirming eating disorder care are made. In this view, this paper offers a foundation for future empirical work in this nascent field of inquiry

by providing a lived experience-led, transdiagnostic, and intersectional account of eating disorders in the context of neurodivergence.

KEY WORDS: neurodiversity, epistemic justice, eating disorders, intersectionality, disability, human rights, gender, autism, adhd

PLAIN ENGLISH SUMMARY

Autistic people and those with attention deficit hyperactivity disorder have a high risk of developing an eating disorder. Other forms of neurodivergence, such as giftedness, intellectual disability, and Tourette's syndrome, have also been associated with disordered eating. Compared to the general eating disorder population, neurodivergent people experience eating disorder symptoms over a longer period of time and commonly experience poorer eating disorder-related treatment outcomes. Yet, there is limited research which is led by neurodivergent people with lived or living experience of eating disorders. However, it is essential to incorporate lived experience expertise in research and treatment design and implementation as this could improve understanding of the many complex and varied underlying eating disorder risk factors which may otherwise guide necessary treatment adaptation(s). Based on the perspectives of lived experience researchers and/or clinicians, this paper discusses the various mechanisms underlying disordered eating and/or eating disorder risk in the context of neurodivergence to inform future developments in eating disorder-related research and clinical practice.

1. POSITIONALITY STATEMENT

All co-authors are neurodivergent and have lived experience of an eating disorder. We are researchers and/or clinicians specialising in the field of eating disorders. We also occupy a wide range of other intersecting marginalised identities across gender, sexuality, body size, class, and neurotype; for example, some of us identify as gender diverse (e.g., transgender, gender non-binary) and/or live in larger bodies. However, we acknowledge that there remain many experiences we cannot speak to as most of us identify as white, none of us has an intellectual disability, and we are all living in Western countries (Australia and the United States of America). Our broad collective personal and professional experiences have informed our investigation of intersectional factors related to disordered eating and/or eating disorder risk, critical appraisal of current eating disorder research and therapies, and considerations for neurodiversity-affirming eating disorder care.

We take on a neurodiversity-affirming theoretical framework. Therefore, we purposefully use non-pathologising and non-stigmatising terminology in relation to neurodivergence. For example, we use terms such as 'traits', 'features', 'characteristics', 'support needs', 'differences', 'atypical', 'challenges', or 'difficulties' rather than 'abnormal', 'symptoms', 'deficits', 'impairments', 'tantrums', 'attention seeking', 'dramatic', 'treatment resistant', or 'non-compliant' [1-3]. We also use identity-first language (i.e., autistic person) in relation to autism as research shows this is the preference of English-speaking autistic individuals [3-4]. Additionally, we refrain from using the terms 'low/high functioning' as these perpetuate erroneous and monolithic stereotypes – that is, autism being a linear construct [5-6] – and fail to capture the dynamic and fluctuating nature of disability [see 7-8].

2. SCOPE AND AIMS

The associations and psychopathological mechanisms underlying eating disorders in the context of neurodivergence remain understudied and poorly understood, as illustrated by limited lived experience-led, transdiagnostic, and intersectional research in this area. With this field of inquiry still in its infancy, this paper does not seek to provide a comprehensive, systematic, review of the literature. Instead, our aim is to provide a narrative review serving as a conceptual, lived experience-led, foundation for future empirical work. In this view, this review builds on the work of Adams et al. [9] investigating the underlying mechanisms linking autism and eating disorders and Field et al.'s [10] exploration of autism-specific eating disorder treatment adaptations. However, given the documented overlapping and transdiagnostic nature of neurodivergence [11-13], our work diverges from Adams et al. [9] and Field et al. [10] as we take a wider lens to investigate neurodivergence using a neurodiversity-affirming theoretical framework. Indeed, our paper presents a holistic overview that considers intersectionality by succinctly capturing the complexity of interconnected socio-cultural and biological factors regarding neurodivergence and eating disorder risk from the perspectives of the authors' broad professional and personal lived experience.

3. BACKGROUND

While evidence demonstrates overall lifetime prevalence of eating disorders (EDs) ranging between 0.74-2.2% for males and 2.58-8.4% for females [14], there are growing concerns over the increasing prevalence of EDs, particularly for children and adolescents [15]. However, pathophysiological mechanisms underlying the development and maintenance of

ED behaviours and cognitions remain poorly understood [16]. Furthermore, less than half of people with EDs who undergo 'evidence-based' treatments make a full, long-term, recovery [17]. The high relapse and mortality rates observed for EDs, even for those who can access treatment [18], have prompted an increased recognition of the need for more lived experience-led research [19] and person-centred care [20-21].

Neurodiversity refers to the diversity of human minds that naturally exists in our society [22]. Neurodivergence refers to individuals who have a neurocognitive profile that diverges from the majority (i.e., atypical), whereas neurotypical people are those whose neurocognitive profile is considered aligned with the majority (i.e., typical) [23]. There are many forms of neurodivergence, such as, but not limited to, autism, attention deficit hyperactivity disorder (ADHD), Tourette's syndrome, giftedness, dyslexia, dyscalculia, dyspraxia, and intellectual disability (ID) [11, 24].

Neuronormativity relies on the assumptions and beliefs which uphold neurotypical privilege—an ideologically oppressive framework which views some personality and neurocognitive traits as inherently superior to and more valuable than others [25-26]. The neurodiversity paradigm, however, rejects the assumption of there being a singular 'normal' personality or neurocognitive profile and opposes the pathology paradigm, whereby those who do not fit within the narrow socially constructed understanding of 'normal' are perceived deficient, disordered, and in dire need of fixing or curing [27-29]. In this view, the neurodiversity paradigm argues that difference should not be regarded as inherently pathological and that the diversity of human minds should be accepted and accommodated for [27, 30].

Research has highlighted high co-occurrence rates between autism, ADHD, and EDs [31-32]. Preliminary research has also associated other forms of neurodivergence, such as ID [33] and

Tourette's syndrome [34-35], with disordered eating. Yet, the mechanisms underlying links between neurodivergence and EDs remain understudied and poorly understood; for instance, there is limited research on the intersectional nature of ED risk factors for neurodivergent individuals (NDI). Subsequently, NDI are shown to experience poorer ED treatment outcomes compared to neurotypical people [36-37, 87].

While there have been calls to include lived experience expertise at the core of mental health research [38], participatory and phenomenological research methods are rarely utilised in the context of EDs [19, 39-41]. With all co-authors being neurodivergent and having lived or living experience of an ED, our narrative review aims to provide a lived experience-based conceptual map to serve as a starting point for future empirical work. A high-level summary of the co-occurring prevalence rates between neurodivergence and EDs is first provided, followed by a thematic investigation of key factors underlying ED risk for NDI. Additionally, a critical appraisal of current ED research and treatments is provided before recommendations for neurodiversity-affirming (NDA) ED care are discussed.

4. METHODS AND CONTRIBUTIONS

A narrative review methodology was selected to provide a broad descriptive framework for this topic, suitable to the overall limited status of literature published to date. In addition, the narrative review is intentionally informed by the lived experiences of the authorship team, all of whom are ED researchers and/or clinicians, providing perspectives and insights with the aim of promoting further conversation and inspiring future research and clinical considerations.

For section five, two databases (PubMed and Google Scholar) were searched (11/03/2024 and 07/06/2024) for prevalence data specific to the co-occurrence discussed (e.g., autism/EDs, ADHD/EDs). Search terms involved a type of neurodivergence (e.g., autism, ADHD, Tourette's syndrome, giftedness) alongside 'eating disorder' or 'feeding disorder'. Specific subsections on prevalence were contributed by authors RE (psychosis and OCD) and LC (autism, ADHD, Tourette's Syndrome, ID, giftedness). The most recent systematic reviews and meta-analyses were selected: here, if no systematic reviews and meta-analyses were identified, the most recent studies were discussed. However, there were instances where no recent studies assessing prevalence were found, in which case it was highlighted that research evidence is limited.

Section six provides a thematic framework whereby factors underlying ED risk in relation to neurodivergence are identified. Co-authors contributed to themes that matched areas of their lived experience, interest, and/or academic expertise; for example, sexuality and gender diversity (authored by SS, with initial outlining by KS), sex diversity (co-authored by SS and KS), giftedness, systemic discrimination, and psychoneuroimmunology (author RE), and sensory processing, executive functioning, systemic discrimination, and psychoneuroimmunology (author LC). Sensory processing and executive functioning differences are discussed first due to their core overarching nature across neurodivergence. Individual characteristics, such as gender identity, sexual orientation, and giftedness follow. Intersectional factors like systemic discrimination and chronic illnesses are then investigated. Section seven consolidates the evidence and positionality to identify current gaps in research

and practice (authored by LC), while section eight provides suggestions for neurodiversity-affirming eating disorder care (co-authored by LC and AR). Furthermore, all co-authors contributed to revision and editing of this narrative review.

5. OVERVIEW OF PREVALENCE DATA

Gillberg [42] was the first to suggest an association between autism and EDs. Throughout the 1990s and early 2000s, sporadic studies have discussed the co-occurrence of EDs and neurodivergence; for example, Fisman et al. [43] found an overlap between obsessive-compulsive traits and anorexia nervosa (AN), Gravestock [44] highlighted an association between ID and binge eating disorder (BED), and Surman et al. [45] posited a link between ADHD and bulimia nervosa (BN). There has been an acceleration of research on this topic in the past decade, consistently reporting high co-occurrence rates between autism and ADHD with various ED diagnoses – namely, AN, BN, BED, and avoidant/restrictive food intake disorder (ARFID) [31-32]. Overall, there is a high degree of overlap and co-occurrence between many forms of neurodivergence [28, 46-47], however, transdiagnostic research regarding forms of neurodivergence other than autism and ADHD in relation to EDs is limited. As such, the evidence below is presented as a high-level summary within the most investigated forms of neurodivergence in relation to EDs – namely autism, ADHD, Tourette's syndrome, obsessive-compulsive disorder (OCD), psychosis, ID, and giftedness.

Despite the increased attention toward EDs in autistic and ADHD populations, there are limitations to estimating co-occurring prevalence rates. Indeed, there is accumulating evidence showing NDI assigned female at birth (AFAB) being erroneously diagnosed with borderline personality disorder [48-51]. One leading argument for such under- and/or misdiagnosis is gender biases in diagnostic instruments and practice [52-56]. Research also suggests that autism and ADHD are underdiagnosed in racially marginalised individuals due to a combination of barriers accessing assessments and ethnocentrism embedded in diagnostic measures [57-58]. Moreover, research shows that autism and ADHD are more

likely to be misdiagnosed as conduct disorder, antisocial personality disorder, and/or oppositional defiant disorder in racially marginalised communities [59-64].

Relatedly, psychometric instruments commonly used for assessing eating psychopathology, as well as body image-related constructs, have not been tailored to, nor validated for, NDI [see 65-66]. Therefore, it is possible that ED measures predominantly used today, such as the Eating Disorder Examination Questionnaire or the Eating Attitudes Test, do not accurately capture EDs and body image disturbances as experienced by NDI [see 65-66]. There is also a paucity of transdiagnostic research investigating neurodivergence and EDs, despite evidenced overlaps between many forms of neurodivergence [46], such as co-occurring autism and ADHD [67], leading to a significant knowledge gap regarding ED presentation and course of illness in these populations.

5.1. Autism

Systematic and meta-analytic reviews have consistently highlighted a strong link between autism and AN: Huke et al. [68] found the prevalence of autism among those with AN varying between 8-37% while Boltri & Sapuppo [69] identified a prevalence range between 8.8-24.5%. Additionally, between 8.2-54.8% of children with ARFID are autistic [31], while 21% of autistic people are deemed at risk of developing ARFID based on a genome-wide association study [70]. Furthermore, undiagnosed ARFID in autistic children represents a risk factor for malnutrition and death [71], highlighting the urgent need for systematic ED screening in autistic children. Between 14-36% of autistic children also show pica behaviours [72-73]. Although research has so far focused mostly on dietary restriction in relation to autism, there is emerging evidence linking autistic traits with all forms of disordered eating, including bingeing behaviours [74-75].

5.2. Attention Deficit Hyperactivity Disorder

There is a significant association between ADHD and EDs. Indeed, in a recent review which included five cross-sectional studies, Villa et al. [32] found that 31.37% of children and adolescent patients with an ADHD diagnosis presented with BED, 19.23% with BN, and 9.38% with AN. Additionally, a meta-analysis found that the prevalence of ADHD in adults with EDs was 3-16.2% for AN, 9-34.9% for BN, and 19.8% for BED [76]. Here, ADHD traits were also found to predict binge eating and bulimic symptoms' severity, even after controlling for anxiety and depression [76]. Moreover, in a study involving 1165 adults diagnosed with an ED, Svedlund et al. [37] found that 37% of individuals with AN and 35% of those with BN scored above the cut-off on the Adult ADHD Self-Report Scale [77]. While prevalence has not been established, research suggests that children with ADHD present an increased risk of loss of control of eating compared to children without ADHD [32, 78]. High incidence of ADHD has also been observed in adolescents [79] and adults [80] seeking weight management surgery.

5.3. Intellectual Disability

There is limited research investigating ID in relation to EDs, in part driven by the lack of adapted ED assessment tools in the context of ID [see 81]. Additionally, there is high heterogeneity, with differing reported ED incidences between people with ID living in the community (1-19%) compared to those who reside in care institutions (3-42%) [44]. Another study showed that 27% of people with ID had an ED, with BED being the most prevalent [82]. In a study of children with ID, Gal, Hardal-Nasser, and Engel-Yeger [33] found that 97% of participants had at least one eating or feeding problem (i.e., food refusal, food selectivity, dysphagia, rumination, binge eating). Furthermore, a systematic review

highlighted a high incidence of swallowing problems (e.g., aspiration, dysphagia) contributing to malnutrition in adults with ID [83].

5.4. Tourette's Syndrome, Obsessive Compulsive Disorder, Psychosis, and Giftedness

There is limited data regarding the co-occurrence of Tourette's syndrome and EDs. However, one study estimated that the lifetime prevalence for EDs was 2% for those with Tourette's, with females (7%) more at risk than males (0.3%) [84]. Furthermore, studies have highlighted a high incidence of disordered eating and feeding challenges for children with Tourette's syndrome [34-35]. For obsessive-compulsive disorder (OCD), meta-analytic prevalence estimates report co-occurrence of OCD in EDs being 13.9 –18%, with greater frequency in AN [85-86]. In comparison to neurotypical individuals experiencing an ED, individuals with OCD or those with elevated OCD traits are at greater risk of more severe ED symptomatology and poorer ED treatment outcomes [87]. Research on the co-occurrence rates of psychotic spectrum disorders and EDs is also limited; however, higher levels of cooccurrence between these two diagnostic groups have been indicated compared to the general population [88]. Indeed, a systematic review showed higher rates of disordered eating compared to controls in individuals with schizophrenia, with disordered eating behaviours observed in 10-41.5% of patients [89]. Moreover, transient psychosis occurs in EDs secondary to malnutrition, electrolyte imbalance, metabolic disturbance, and re-nutrition [90-91] and incidence of transient psychosis among ED patients has been estimated to be 10-15% [91]. Lastly, some studies have suggested a link between giftedness and EDs, with gifted individuals thought to be at a high risk of developing EDs; however, there is no prevalence data available [92-98].

5.5. Conclusion

While clear associations between autism, ADHD, and EDs have been highlighted, more investigations are needed with regards to other forms of neurodivergence. Additionally, despite many forms of neurodivergence overlapping and co-occurring, there is limited transdiagnostic research investigating neurodivergence and EDs. Nonetheless, estimating prevalence remains challenging due to issues such as NDI being misdiagnosed with personality disorders and lack of ED-focused psychometric instruments tailored to, and validated for, NDI.

6. NEURODIVERGENCE, INTERSECTIONALITY, AND EATING DISORDERS

The reasons underlying NDI being at a high risk of disordered eating and/or EDs are varied, complex, and multifaceted, and encompass a wide range of interconnected environmental, psychosocial, and biological factors. NDI experience the world differently in a myriad of ways, as illustrated by differences in sensory processing, communication and socialising (e.g., the double empathy problem), thinking and learning styles (e.g., monotropism, inertia), gender identity, sexual orientation, sex diversity, and executive functioning [99-101].

Together, these differences may contribute, either directly or indirectly, to a wide array of pathways potentially contributing to the development of disordered eating, EDs, and/or body image disturbances for NDI [9, 19, 101-103].

The section below outlines several common neurodivergent experiences that may contribute to elevated ED risk. The available evidence has been integrated with the lived and living

experiences of the authorship team to provide a framework for future work on this emerging topic within the ED field.

6.1. Sensory Processing

Sensory processing differences encompassing exteroception (related to external stimuli such as sounds or tastes), interoception (related to internal stimuli such as thirst, pain, or hunger), and alexithymia (the ability to recognise and/or describe affective states and associated physiological sensations) are core overarching characteristics associated with neurodivergence [104-112]. Collectively, atypical sensory processing patterns are linked to EDs as well as body image disturbances, irrespective of neurotype [113-115]. Additionally, research shows that atypical sensory processing, as experienced by autistic people more specifically, is strongly associated with disordered eating [116-118].

Exteroception and interoception influence cognitive processes such as goal-directed behaviours (e.g., eating, drinking), motor control and coordination, memory, learning, motivation, self-perception, and emotional awareness and responsiveness, and are tied to neurobiological systems, particularly thalamo-cortical connectivity [119-121]. The thalamus is located at the centre of the brain and has historically been perceived as a passive relay centre for sensory information, including interoception and exteroception (except for olfaction, which is routed through the olfactory bulb) [122]. However, a growing body of evidence shows that the thalamus plays a more active and important role in many cognitive functions than previously thought, including regarding memory and attentional control [123-124], executive functioning [125], and emotional processing (e.g., thalamo-limbic connectivity) [126]. More recently, neuroscience research has suggested the thalamus to be involved in a wide range of psychiatric conditions (e.g., depression, anxiety, bipolar disorder,

post-traumatic stress disorder, substance abuse) [127-129], including EDs [130-132]. Atypical sensory processing likely underlies differences in eating processes and habits as well as vulnerability to ED development [see 133, pp. 22-23].

There are different ways in which atypical sensory processing may lead to eating difficulties and/or disorders. Exteroception may contribute to dietary restriction resulting from aversions whereas sensory seeking and associated cravings tied to specific tastes or textures may underlie bingeing [113]. Heightened exteroception (e.g., hyperacusis, misophonia) may also be contributing to the preference for eating alone to avoid sensory overwhelm [134-135]. Moreover, interoception may impact eating through the modulation of hunger and satiety; for instance, feeling full after consuming small amounts of food may underlie restriction whereas delayed satiety may promote bingeing and/or eating in the absence of hunger [113]. Interoception further plays an important role in body image and has been associated with body image disturbances [102, 136-137]. Research has also associated alexithymia, which is connected to interoception, with binge eating behaviours (often referred to as 'emotional eating) [138-139]. Specifically, Vuillier et al. [140] found that alexithymia could partially explain the high prevalence of EDs in autistic people.

Eating a restricted range of foods is often framed as an ED symptom. While this phenomenon may reflect psychopathology for some, qualitative research shows that, for autistic people, it may be an adaptive and self-regulatory mechanism to prevent sensory distress, cognitive overwhelm, and mealtime anxiety [103, 141-142]. Indeed, research suggests that autistic people do not habituate to sensory stimuli in the same ways as non-autistic people [143-144]. In the context of such reduced habituation, coercing autistic people to eat aversive foods may be experienced as traumatic and unintentionally exacerbate feeding difficulties (e.g., food phobia) [145]. It is therefore critical to take the different ways NDI experience eating and

feeding from a sensory perspective into account for ED identification, care, and recovery [9, 10, 19, 36].

6.2. Executive Functioning

Executive functioning differences (e.g., task initiation, memory, demand avoidance) are influenced by sensory processing patterns [146] and are observed in many NDI [147]. Executive functioning differences are also linked to EDs [148-149]; for example, executive functioning differences may contribute to difficulties related to remembering to eat, grocery shopping, meal preparation, consistently following a meal plan, and/or choosing what to eat [113, 101, 141]. Such difficulties may promote restrict-binge cycles, whereby forgetting to eat or being unable to organise meal preparation leads to prolonged periods of time without eating, which may be followed by compensatory bingeing episodes [101]. Additionally, given that many autistic people have a literal, or concrete, thinking style [23, 150], dichotomous portrayals of 'healthy' or 'good' versus 'unhealthy' or 'bad' foods associated with diet culture and public health messages may contribute to fixations towards foods considered 'healthy' (e.g., orthorexia) [101, 151-152].

6.3. Gender, Sexuality, and Sex-Diversity

People with a diverse sex, sexuality, and/or gender identity are more likely to be autistic and/or have ADHD [153-159] and face a high risk of EDs and body image disturbances [160-161]. Factors contributing to the increased risk of EDs among neurotypical sex, sexuality, and gender diverse people include gender dysphoria, internalised cis-heterosexist ideals, and minority stress [133, 160, 162-168]. In contrast, identity pride, community connectedness, and gender-affirming healthcare can be protective factors [133, 160, 166, 169].

Trans people experience disproportionate rates of EDs compared to cisgender individuals [162-164, 170]. EDs, in this context, may function as a way to manage gender dysphoria; for instance, restrictive eating can delay the onset of puberty or minimise the appearance of secondary sex characteristics [162-164]. ED risk for trans people is further exacerbated by minority stressors, including discrimination and social exclusion, which can lead to heightened body dissatisfaction and increased engagement in ED behaviours [171]. Additionally, high levels of gender dysphoria are observed in the autistic population, partially mediated by atypical sensory processing, contributing to greater difficulties in coping with gender incongruence and body image disturbances [157, 172-174].

Variations of sex characteristics (VSC) – otherwise referred to as intersex— are differences in innate sex characteristics that differ from social and medical norms [175]. In a study of 1022 people with VSC, 8.8% had a diagnosis of autism, 9.1% had high autistic traits, and 4.1% screened positive for ADHD [155]. Higher autistic trait scores were associated with lower self-esteem in participants with Turner Syndrome, Klinefelter Syndrome, and female participants with 46XY variations with and without androgenisation [155]. Additionally, higher ADHD trait scores were associated with poorer body image in participants with Turner Syndrome, female participants 46XY variations with and without androgenisation, and male participants with 46XY variations [155]. ED diagnoses were reported in 11.1% of the overall sample (3.7 times higher than the general population); however, stratified data of ED diagnoses for autistic intersex people and those with ADHD were not provided [155]. Intersectionality plays a crucial role in understanding elevated ED risk for NDI who are also members of the LGBTQIA+ community, highlighting the need for targeted interventions and support systems that address the unique intersections of neurodivergent and LGBTQIA+ identities [153, 161]; for example, by acknowledging that dietary restriction may stem from

different motivations in this population (e.g., sensory and gender dysphoria-based), underscoring the importance of neurodiversity-affirming ED care also being gender-affirming.

6.4. Giftedness and Twice-Exceptionality

Giftedness, which refers to outstanding intellectual and/or creative potential [176], may represent a risk for ED development through perfectionism, low self-esteem, striving to achieve social acceptance, and avoiding ostracization and rejection [177-179]. Obsessive perfectionism may develop in gifted people within the context of interpersonal trauma and marginalisation, wherein an individual pursues creative and/or intellectual achievements as a survival process to protect autonomy, identity and individuality, and to anesthetise psychological pain [180-182]. High academic and/or creative achievement in the context of giftedness may be an asset to accepting uniqueness and social difference; however, this may become a burden when an individual's emotional vulnerability and associated need for support are not recognised and addressed [183]. Twice-exceptional people (i.e., people who are both disabled and gifted, such as those with co-occurring giftedness and dyslexia and/or ADHD) may also experience psychosocial vulnerability and imposter syndrome due to asynchrony – referring to disparities between ones' chronological and developmental trajectories, belongingness with peers, and expectations [177]. Perfectionism and disordered eating may be an attempt to cope with the psychosocial distress [see 184] associated with such asynchrony [97, 177, 185]. Twice-exceptionality may also present a social and identity threat, due to experience of being 'twice different' [186]; thereby, disordered eating may serve as a social survival strategy to fit in with peers through striving toward societal body ideals [177] and alleviate the existential angst that may accompany fears of failure and/or imposter syndrome [187].

6.5. Systemic Discrimination, Inequity, and Trauma

NDI commonly experience systemic oppression and discrimination across varying settings and contexts (e.g., education, healthcare, employment) [188-195]. Such forms of systemic discrimination increase the likelihood of unemployment [196], houselessness [197], and food and nutrition insecurity [198-199], therefore contributing to psychosocial distress and mental ill-health [200], including EDs [201]. Indeed, high rates of houselessness and unemployment have been reported for NDI [196, 198, 202-205]. The risk for autistic people to be food insecure is four times greater compared to non-autistic people [198]. Moreover, a systematic review highlighted that between 10-40% of adults with ID experienced houselessness [206]. Food insecurity is a known ED risk factor, particularly for BN [207-209].

Rejection and stigmatisation are common experiences for NDI, and negatively impact on self-esteem and psychosocial wellbeing [210-212]. Rejection and stigmatisation are often rooted in deficit-based and dehumanising stereotypes, such as that autistic people 'lack empathy' and 'social skills' [213-216] or that those with ADHD are 'lazy' and 'attention seeking' [217-218]. Resultantly, NDI are prone to internalise such stigma and may hide their authentic selves to avoid being rejected and bullied – a phenomenon referred to as *masking* or *camouflaging* [219]. Masking is associated with loss of sense of self and identity confusion [220], which are known risk factors for EDs [221]. Furthermore, substantive evidence supports the link between masking and poor mental health outcomes in autistic people, such as anxiety, self-harm, suicide [222-225], and risk of EDs [151, 226].

NDI have been found to experience higher rates of interpersonal victimisation, including domestic abuse and sexual violence, compared to neurotypical individuals [227-231]. For example, nine out of ten autistic women [232] or those with an ID [233] report having been

sexually assaulted, which is three times more than neurotypical women. Additionally, a systematic review suggested that those with ADHD were more likely to have experienced childhood sexual abuse compared to those without ADHD [234]. Such experiences of victimisation are relevant as interpersonal trauma, including sexual abuse, have been found to substantially increase the risk of EDs [201, 235-237].

There is emerging evidence suggesting that autistic people experience post-traumatic stress triggered by factors not currently conceptualised within psychiatric nosology as inherently traumatic [238-241]; for example, some autistic people may experience post-traumatic stress resulting from repeated exposures to aversive sensory input (e.g., sounds, smells, tastes, textures) [145]. It is estimated that up to 60% of autistic people suffer from post-traumatic stress disorder in their lifetime [241], with many thought to be misdiagnosed (e.g., borderline personality disorder) and not receiving the trauma-informed care they need [49-50, 239]. Since post-traumatic stress disorder is a known contributor to the development of EDs [242], the delayed identification and delivery of compassionate treatment of trauma may therefore exacerbate the vulnerability towards developing EDs for autistic people.

Intersecting experiences of marginalisation and structural oppression (i.e., racism and ethnocentrism, ableism, gender binarism, sexism, cis-heterosexism, and sizeism) may compound ED risk [237, 243-244]. Indeed, in Nelson et al.'s [237] longitudinal study involving 331 undergraduate students, different forms of discrimination such as racism, weight stigma, and sexism all predicted body image dissatisfaction and disordered eating. The patterns of disordered eating were different depending on the type(s) of discrimination experienced [237], highlighting the importance of considering the influence of intersecting factors of identity, oppressive power structures, and trauma experiences in ED development and recovery, irrespective of neurotype [20].

6.6. Chronic Illness and Psychoneuroimmunology

NDI experience chronic medical illnesses at higher rates than neurotypicals [245-247]. The range of such medical conditions include, but are not limited to, Ehlers-Danlos syndrome, postural tachycardia syndrome [248], diabetes type 2 [249], polycystic ovarian syndrome [250], endometriosis [251], asthma, dermatitis, food allergies [252-253], gut microbiome dysbiosis [254-255], and gastrointestinal disorders [256-257]. There is emerging evidence suggesting a bidirectional relationship between the aforementioned conditions and EDs; indeed, experiencing one of the conditions listed above may increase the risk of developing an ED later in life, while having an ED may increase the risk of developing one of the aforementioned conditions later in life [258-263].

Furthermore, there is growing evidence from the field of psychoneuroimmunology suggesting that psychological stressors, including in relation to systemic discrimination, interpersonal victimisation, and childhood trauma, may be a key factor in immune and gutbrain dysregulation [264-267]. It has been suggested that the link between psychological stress and gut-brain and immune dysregulation is rooted in the mechanism whereby stress activates corticotropin releasing hormone (CRH) which, in turn, triggers the degranulation of mast cells [268-270]. More specifically, research points to a bidirectional relationship between the immune system, particularly mast cells and histamine [271-272], and a wide range of psychiatric disorders [273-274], including EDs [262, 275-277]. Some research also posits that this relationship may be mediated by histamine playing a role in the modulation of dopamine neurons' firing rates in the brain [278], particularly in the thalamic region [279-280]. Histamine further influences glucose and lipid metabolism [281], leptin and ghrelin activity (i.e., appetite, fullness) [282], digestion [283], and control of arousal and energy expenditure (i.e., homeostasis), including propensity towards exercising [284-286]. As noted

by Keeler et al. [287], "infections, autoimmune diseases, and stress-related alterations in the immune system influence brain circuits and neurotransmitters involved in appetite and mood regulation".

6.7. Conclusion

There are many intersectional factors underlying NDI being at a high risk of experiencing disordered eating and/or developing EDs. This underscores the importance of investigating ED risk through a holistic, intersectional, and lived experience-led lens. Taking these intertwined factors into account in treatment formulation is also of great importance to improve ED care individuation.

7. GAPS IN RESEARCH AND PRACTICE

EDs are psychiatric disorders with significant negative consequences on psychosocial wellbeing and physical health [288]. Despite this, our current understanding of mechanisms for ED risk, development, maintenance, and relapse remain poorly understood, irrespective of neurotype [18, 289].

While manualised psycho-behavioural mental health approaches, such as cognitive-behavioural therapy (CBT), are often promoted as first-line 'evidence-based' treatments for most mental illnesses, including EDs [17-18], there are a range of issues that impact the generalisability of such broad claims [290]. Identified issues include publication bias [291-293], selective reporting [294-295], undisclosed conflicts of interest [296-297, see 298], under-reporting of adverse effects [299-301], lack of long-term follow-up studies [18, 302], and lack of participatory, lived experience-led, research [39, 41, 303].

Indeed, psycho-behavioural treatments considered as the 'gold standard' in the context of EDs – namely, CBT (for adults) and family-based therapy (FBT; for young people) – have consistently been shown to lead to less than half of people with EDs reaching long-term recovery overall [17-18, 304]. There is also emergent evidence showing that NDI, more specifically, have poorer ED treatment experiences and outcomes compared to neurotypical individuals [87, 305-310].

7.1. Reporting of adverse effects and risk assessments

Insight into both intervention efficacy and risk of harm is essential to the practice of evidence-based medicine; indeed, as noted by Junqueira et al. [311], "treatment decisions rely on accurate knowledge of both efficacy and harms". However, adverse effects related to psycho-behavioural treatments for mental ill-health frequently lack systematic assessment and transparent reporting in research, subsequently skewing evidence regarding risk-benefit ratio [299-300, 312-315]. For example, in a systematic meta-review of meta-analyses of ED treatments, adverse events were only acknowledged in the context of pharmacology [316]. Furthermore, Linardon et al. [317] conducted a systematic review and meta-analysis to assess the efficacy of CBT for treating EDs where they deemed CBT to be efficacious despite omitting to discuss measures of short and long-term harms. Iatrogenesis, that is, harm resulting from healthcare practices, may not manifest immediately and be delayed [299], in which case it would only be captured in long-term follow up studies. Late onset iatrogenic harm is therefore missed by the lack of long-term follow-up studies in ED research [see 18]. Subsequently, lack of reported iatrogenic harm in the literature is mistakenly interpreted as evidence of no harm [see 301], leading mental health practitioners under-resourced to recognise and address iatrogenesis from psycho-behavioural treatments [see 300, 318], including in the context of neurodivergence [36, 319-321].

There is an increasing number of community resources where NDI with lived experience of EDs report having experienced iatrogenic harm from ED treatments, including psychological trauma [322-326]. Unfortunately, such narratives are seldom reflected in published mental health clinical trials and meta-analyses. Biases undervaluing qualitative, phenomenological, and lived experience-led research have been suggested to contribute to such narratives not being adequately discussed in mental health research [327-328]. Nonetheless, there is growing recognition that psycho-behavioural treatments can be harmful to NDI [318-319, 329], including in the context of EDs [19, 309, 330-331].

7.2. Neurodivergent Leadership and Representation: Who is Creating The 'Evidence' and Who is the 'Evidence' For?

According to Curnow et al.'s [332] umbrella review investigating the overall suitability of currently available mental health therapies in the context of autism, no existing mental health framework can claim to be NDA; a conclusion that has also be highlighted in the ED context [10, 36, 309]. There are two main factors, discussed below, playing into this: namely, the lack of representation of NDI in mental health research and the lack of non-tokenistic, lived experience-led, participatory research. Indeed, "inclusive research practice characterized by the involvement of neurodivergent people in meaningful roles has been described as a 'requirement of excellence' in neurodiversity research" [333].

Participatory research methods which are led by individuals with lived experience are rarely utilised in ED research [39-41] or in mental health research more broadly [334-337, pp. 179-180]. It has been suggested that participatory research methods have potential to improve overall mental health outcomes for NDI [336, 338-340], including in relation to EDs [19, 40]. Indeed, participatory research methods have been shown to facilitate the translation of

findings into real-world contexts and ensure that the research yields relevant and meaningful benefits for the community [340].

The neurodivergent community, and other marginalised communities (e.g., gender diverse folks) [341], are not adequately represented in ED research [10, 19, 40]. The lack of neurodivergent representation is not limited to EDs; for example, Jubenville-Wood et al. [303] noted that "limited research exists that examines the experiences of autistic individuals in psychotherapy." The neglect of intersectionality in ED research further extends to co-occurring mental illnesses, despite a majority of those with an ED also being diagnosed with another psychiatric condition [342]. This is particularly relevant in the context of neurodivergence as NDI are shown to experience mental illnesses, including other than EDs, at significantly higher rates compared to neurotypical individuals [343-347].

A greater focus on participatory research that takes an intersectional lens is indicated to better understand the connections between neurodivergence and EDs and improve ED prevention efforts and treatment outcomes for NDI [10, 40, 174, 264, 331].

7.3. One Size Does Not Fit All

The indiscriminate use of a narrow range of manualised therapies, assessment tools, and public health campaigns for different demographic groups, such as NDI, gender and/or sexually diverse people, and racially marginalised individuals, is problematic, as each group experiences unique underlying factors to the development of EDs and may require different, sometimes conflicting or opposing, forms of support and formulations to recover [see 10, 174, 348-349]. As outlined in section six, NDI, as well as those with other minority and/or marginalised identities, have been found to have divergent feeding, eating, and ED experiences. However, hegemonic conceptualisations of feeding, eating, and EDs do not

incorporate such diversity of lived experiences, a limitation that impacts all stages of ED care [see 10, 174, 348-349].

Evidence-based medicine requires that clinicians consider research evidence, clinical expertise, and patient preferences when deciding on a course of action [350]. Indeed, ED clinical guidelines identify a narrow range of 'evidence-based' ED interventions, many of which are manualised, whilst also acknowledging the important role of person-centred approaches to ED care [351-352]. This is a realistic goal when the manualised treatments are informed by diverse and representative samples that adequately capture divergent experiences and account for them. However, given the traditional misrepresentation of authentic and inherently adaptive neurodivergent eating and feeding experiences as pathological, and the historical and ongoing legacy of epistemic injustice silencing neurodivergent (and other marginalised) voices in ED research [see 19, 40, 164, 174, 348-349], it appears that oppressive and unequal power dynamics have contributed to the lack of more affirming, innovative, and inclusive conceptualisations of and interventions for EDs, setting the scene for the monopolisation of ED research by a limited number of manualised interventions (e.g., CBT, FBT), despite their lacklustre success rates [see 17-18, 304].

Importantly, this legacy of epistemic injustice places ED clinicians who are attempting to translate research evidence into clinical practice in a double bind: on the one hand, clinicians must practice in accordance with ED clinical guidelines which identify a narrow range of suitable ED treatment approaches, typically manualised, that they can choose from, whilst simultaneously being asked to practice person-centred care that, if followed, is not likely to fit neatly into the prescribed formulation and treatment plan laid out in the approved manualised interventions.

The RAVES model (an acronym representing regular eating, adequacy, variety, enjoyment, and social eating) is a clinical tool that is commonly used by dietitians [353]. Whilst the intervention targets identified in this clinical tool appear logical when framing eating through a neuronormative lens, they do not align with lived-experience informed conceptualisations of adaptive neurodivergent feeding and eating experiences, such as preference for sameness [142], sensory-related food variety constraints [see 9, 40, 151], and preference to eat alone [134], and risks invalidating authentic and adaptive neurodivergent feeding and eating support needs and preferences if applied indiscriminately [101].

Food-related exposure-based techniques, typically used in tandem with psycho-behavioural ED treatments such as FBT and CBT [see 354], risk causing sensory overwhelm, distress, or even sensory trauma when they fail to account for and accommodate divergent sensory processing profiles [see 142, 145], particularly in light of the evidence showing that sensory habituation is attenuated amongst autistic individuals [143, 355-356].

Research demonstrates that those with atypical AN experience similar psychopathology, illness severity, and medical risks to those with typical AN, regardless of weight status [357-359]. The current weight-centric conceptualisation of AN recovery is thus problematic, particularly for autistic people who report that weight is not the most predominant factor contributing to their food restriction [9, 10, 141, 151, 174, 309]. Indeed, research investigating treatments for AN, including CBT, mainly focuses on weight gain as a main indicator of recovery and a primary outcome measure for treatment effectiveness [36-361], despite growing evidence that weight does not measure recovery and that focusing on weight in this manner contributes to weight stigma (e.g., being turned away from health services due to being perceived as 'not being sick enough') [362-364] and iatrogenic harm [365].

emphasise the importance of supportive interpersonal connections, strengths-based sense of identity, finding meaning and purpose in life, and orienting towards self-compassion as central factors supporting ED recovery, more so than weight [21].

7.4. Conclusion

Current psycho-behavioural frameworks for treating EDs, such as FBT and CBT, are often broadly promoted as effective and 'evidence-based', despite meta-analytic evidence consistently demonstrating only less than half of people who access these treatments experience long-term recovery, irrespective of neurotype. Additionally, evidence shows that autistic people and those with ADHD experience poorer treatment outcomes compared to neurotypical people; unfortunately, there is a paucity of empirical evidence for other types of neurodivergence. Issues relating to lack of long-term follow-up data, limited assessment and reporting of iatrogenesis, undisclosed conflicts of interest, and protocol and publication biases, have been identified as factors to consider when interpreting efficacy claims in this context. More specifically, participatory research methods are seldom used in ED research and NDI are rarely represented or taken into consideration in treatments' design and implementation, as are other marginalised communities, thus significantly undermining the validity of 'evidence-based' mental healthcare for a more diverse range of people. Furthermore, ED research rarely takes a holistic lens that acknowledges intersectional factors, as outlined in section six, needing to be considered in treatment formulations, leading to gross oversimplifications in manualised treatment frameworks that do not account for the fact that different people and communities have different, and sometimes conflicting, support needs.

8. WHAT DOES NEURODIVERSITY-AFFIRMING EATING DISORDER CARE ENTAIL?

The concept of NDA care is grounded in the principles of the neurodiversity paradigm which views neurodiversity as a natural and valuable form of human diversity, rejecting the idea that there is a singular 'normal' personality and neurocognitive profile that all humans should universally aspire to and emulate [22]. An NDA mental healthcare framework, therefore, is a holistic and person-centred framework rooted in the principles of phenomenology, epistemic justice, and cultural humility [339, 366-371]. NDA ED care requires that ED clinicians engage in ongoing self-reflection to challenge biases and neuronormative assumptions, social norms, beliefs, and goals that further perpetuate the pathologisation and systemic oppression of NDI [40, 101, 174, 264]. Crucially, NDA ED care necessitates clinicians be able to differentiate neurodivergent traits from ED symptoms, understand the underlying reasons and adaptive mechanisms of neurodivergent traits, and validate and accommodate the individualised support needs of NDI, such as with regards to sensory processing, communication, and executive functioning, rather than attempt to 'fix' these at the expense of overall wellbeing and self-determination [see 40, 101, 174, 264, 372].

8.1. Overview of currently available neurodivergent-specific psychometric assessments and treatment adaptations

Specific assessment tools have been developed to improve the identification of feeding and eating difficulties, and related psychopathology, amongst autistic individuals (Swedish Eating Assessment for Autism spectrum disorders [373]; Body Appreciation Scale 2 for Autistic Adults [65]; Autism Eating Questionnaire [374]; Brief Autism Mealtime Inventory Assessment [375-376]. To the authors' knowledge, no assessment tools have been developed

or adapted to target eating or body image psychopathology associated with other forms of neurodivergence.

Suggestions have also been made to tailor treatments for NDI [10, 377-378]. However, except for the Body Appreciation Scale 2 for Autistic Adults [65], none of the adaptations to either identification or treatment discussed in this section have been explicitly identified by their authors as having been developed through lived experience-led research. It is also important to note that while care adaptations discussed below have been developed with the intention of improving treatment outcomes for NDI, this does not necessarily guarantee that they will be experienced as NDA by NDI with EDs or helpful in practice [see 379]. Empirical evidence led by NDI with ED lived experience is required to provide insight on whether these ED care adaptations are experienced as helpful, harmful, or neutral.

An ED inpatient care pathway tailored for autistic people, the PEACE Pathway, has been implemented in the United Kingdom [377]. The PEACE Pathway outlines the need for individualised sensory processing accommodations and includes training for clinicians to assist in making autism-specific adjustments to ED care and psychoeducation [380]. A preliminary evaluation of the PEACE pathway found that it reduced treatment duration and resulted in admission cost savings [380]. The PEACE pathway researchers examined the feasibility of implementing Cognitive Remediation Therapy (CRT) amongst individuals with an ED, comparing efficacy between those with low and high autistic traits, and found CRT supported improvements in cognitive flexibility and central coherence in both groups [381]. However, across studies, results are heterogeneous, with Saure et al.'s [115] narrative review concluding that individuals with AN with high autistic traits appear to benefit less from CRT (as well as from cognitive remediation and emotion skills training interventions) compared to those with low autistic traits.

Drawing on available research and clinical experience of supporting autistic children and young people with AN, Loomes and Bryant-Waugh [382] identified a range of autism-specific modifications for family-based interventions for AN. The suggested adaptations target four domains: sensory processing considerations, cognitive and behavioural factors, difficulties with social communication and relationships, and challenges relating to emotion processing [382]. The authors recognise that the suggested adaptations are conceptual and thus require empirical assessment before we can determine their acceptability and efficacy.

Field and colleagues [10] undertook a similar project, identifying ways in which ED services could better meet the needs of autistic women via a Delphi Study. Fifty-six consensus statements reached agreement in the final round, relating to the need to support the unique experiences of autistic women, separating autistic features from the ED, and the importance of staff being experienced in supporting autistic individuals [10].

A single case report detailed adaptations to CBT to meet the needs of an individual with ID experiencing difficulties with binge eating [383]. Adaptations included a focus on regularity and predictability of appointments, the use of images to aid with understanding, adapted use of language, and sensory considerations. The authors reported that the participant's self-regulation associated with loss of control eating improved as a result of the intervention [383].

8.2. Human Rights, Cultural Humility, and Epistemic Justice

An NDA approach to mental healthcare is a human rights-based approach rooted in the social model of disability and existential-humanistic psychology [339, 366-368, 384]. As such, NDA mental healthcare is inherently gender-affirming and trauma-informed [339, 366]. This requires the rejection of compliance and exposure-based behavioural approaches (e.g.,

operant conditioning, contingency management) that are imposed rather than self-determined and client-led as these have been associated with iatrogenesis, including learned helplessness and psychological trauma [225, 339, 385-389]. Instead, the priority must be on supporting self-determination, self-advocacy, and self-efficacy, and incorporating a person-centred, compassionate, and strengths-based approach to care that is informed by lived experience expertise [19, 337, pp. 179-180, 366, 390-393].

It is imperative that all NDA mental health interventions aim to support the development of a positive and authentic neurodivergent identity, as this has been associated with benefits to overall psychosocial wellbeing and increased self-compassion [394-395]. The benefits of a positive neurodivergent identity have also been noted in relation to supporting ED recovery specifically [10, 19, 151, 174]. Such an approach echoes the recent call to action released in a joint statement by the United Nation and the World Health Organization that urges mental health practitioners and researchers to focus on the social determinants of mental health, epistemic justice, harm reduction, and human rights-based mental healthcare [396].

8.3. Reasonable Adjustments

Founding members of Autistic Doctors International have created SPACE; an autistic-led framework for autism-specific accommodations aimed at improving accessibility of healthcare services for autistic individuals [370]. The SPACE framework identifies key reasonable adjustments across core support needs for autistic people, such as sensory, predictability, acceptance, communication, and empathy, as well as across three main domains of experience: physical space, processing space, and emotional space [370]. As noted in section six, NDI have unique sensory, communication, cognitive, and emotional needs that have been identified as important to understand, validate, and accommodate as part

of NDA ED care, and as such, SPACE has potential utility as a tool to support more neurodiversity-affirming ED service delivery [see 9, 10, 101]. For example, some autistic people may be non-speaking, while others may be partially speaking (e.g., situational mutism), and allowing communication in ways that feel comfortable is likely to improve trust between patient and clinician [370]. Additionally, adapting the consulting room to meet the sensory needs of autistic patients may also assist in building trust, self-regulation, and safety (e.g., noise, temperature, smells, lights) [370]. On the other hand, some autistic patients may prefer to remain standing or sit on the floor during the consultation, and these differences in proprioception should also be accommodated [370].

Research exploring general mental health has highlighted empirical evidence supporting the benefits of sensory-based approaches in the general population, where the physical environment is tailored to the sensory processing characteristics of an individual [397-401]. Sensory-based approaches can include strategies such as adjusting the room's lighting and/or colours, providing access to weighted blankets, fidget tools, rocking chairs, and/or noise-cancelling headsets to reduce sensory stimulation. Sensory-based mental health approaches have been shown to create a sense of safety and control, increase self-regulation, reduce distress and anxiety, improve self-perception, and help with stabilising (or 'grounding') acute arousal [399, 401]. There is also emerging evidence suggesting that sensory-based approaches in inpatient mental health facilities may contribute to improvements in autistic individuals' wellbeing and self-regulation [372, 402-403], giving further weight to the argument in support of ED services attending to the unique sensory needs of NDI.

8.4. Conclusion

NDA care involves respecting and validating different personality and neurocognitive profiles instead of de facto pathologising divergence and trying to 'fix' differences rather than celebrating diversity and accommodating specific support needs. Promoting self-determination and a positive, authentic, sense of self are critical to NDA care. In the context of EDs, NDA care requires that researchers and clinicians engage in epistemic justice and remain curious towards the different ways NDI engage with eating, self-perception and expression, and conceptualise recovery. Indeed, distinguishing between ED symptoms and adaptive ND traits is key to NDA ED care and the prevention of iatrogenesis. Additionally, while further empirical research is required, the SPACE framework and sensory-based approaches may be useful in the context of ED care for NDI.

9. CONCLUSIONS

Prevalence data highlights that autistic people and those with ADHD are at an increased risk of developing EDs compared to those who are neurotypical. The limited research on other forms of neurodivergence (e.g., giftedness, intellectual disability, Tourette's syndrome) and eating psychopathology has found them to be associated with high incidences of disordered eating, highlighting a need for more research. It is possible that the real co-occurrence rates between neurodivergence and EDs are much higher as issues such as the underdiagnosis and/or misdiagnosis of autism and ADHD in AFABs and racially marginalised populations, lack of tailored psychometric instruments, and limited research despite documented overlaps between various forms of neurodivergence potentially lead to under-identification of true prevalence rates. Improvements in the accessibility of neurodivergent assessments and the development of lived experience-led ED screening and diagnostic tools are likely to support

early identification for NDI. In addition, given the overrepresentation of neurodivergence in ED populations, it is ideal for ED clinicians to screen for autism and ADHD to improve ED care individuation.

There are a wide range of interwoven factors underlying ED risk for NDI, ranging from atypical sensory processing, executive functioning, social communication, emotional processing, and a higher likelihood of experiencing chronic illnesses and systemic discrimination. Therefore, it is important that future investigations into ED risk in the context of neurodivergence take an intersectional approach. Transdiagnostic research factoring in the overlaps between varied forms of neurodivergence is also warranted given that co-occurring forms of neurodivergence appear to have an impact on ED presentations and the nature of support needs. Future empirical work should be participatory, non-tokenistic, and ideally lived experience-led, and acknowledge power structures associated with discrimination and trauma to better achieve epistemic justice and better translate to cultural humility in ED care.

10. STRENGTHS AND LIMITATIONS

While our work did not entail a comprehensive systematic review of the literature due to the nascent nature of this specific area of inquiry, this narrative review allows for scoping of the literature available to date whilst providing a lived experience-led framework for discussion and an initial conceptual map for future empirical work.

Additionally, this review can only draw from the strength of the evidence that is available.

Although an extensive narrative review, research is still limited, and this can only be improved with funding and support. Furthermore, most of the literature to date focuses on

NEURODIVERGENCE, INTERSECTIONALITY, AND EATING DISORDERS: A LIVED

EXPERIENCE-LED NARRATIVE REVIEW [PREPRINT]

either autism or ADHD in relation to EDs. Therefore, there is limited data relevant to other

forms of neurodivergence to draw upon.

This paper was written exclusively by NDI with lived or living experience of an ED, which

may influence reflexivity. However, the importance of lived experience-led mental health

research has been consistently highlighted, particularly with regards to translation and

implementation.

LIST OF ABBREVIATIONS

ADHD: attention deficit/hyperactivity disorder

AFAB: assigned female at birth

AMAB: assigned male at birth

AN: anorexia nervosa

ARFID: avoidant/restrictive food intake disorder

BN: bulimia nervosa

BED: binge eating disorder

CBT: cognitive behavioural therapy

CRH: corticotropin releasing hormone

CRT: cognitive remediation therapy

ED: eating disorder; EDs: eating disorders

FBT: family-based therapy

ID: intellectual disability

NDI: neurodivergent individuals

36

NEURODIVERGENCE, INTERSECTIONALITY, AND EATING DISORDERS: A LIVED

EXPERIENCE-LED NARRATIVE REVIEW [PREPRINT]

OCD: obsessive compulsive disorder

VSC: variations of sex characteristics

DECLARATIONS

Ethics approval and consent to participate:

N/A

Consent for publication:

N/A

Availability of data and materials:

All data generated or analysed during this study are included in this published article.

Competing interests:

Laurence Cobbaert, Anna Rose, and Elysia Thomas are on the Board of Directors of not-forprofit organisation Eating Disorders Neurodiversity Australia (EDNA). They receive no direct financial remuneration from their Directorship positions with EDNA.

Laurence Cobbaert is the Founder and Chair of the Neurodiversity special interest group at the Australia and New Zealand Academy for Eating Disorders (ANZAED), and a member of the Australasian Society for Autism Research (ASfAR).

Anna Rose has received financial contributions in relation to their lived experience consultation for the Australian Eating Disorders Research and Translation Centre (AEDRTC), and professional development consultation for the Australia and New Zealand

37

Academy for Eating Disorders (ANZAED). Anna also receives a PhD stipend from Bond University, Queensland, Australia.

Funding:

This paper is partially inspired by the technical report titled *Eating Disorders and*Neurodivergence: A Stepped Care Approach (March 2023), which was commissioned by the National Eating Disorders Collaboration (NEDC) and co-authored by Laurence Cobbaert and Anna Rose on behalf of the not-for-profit organisation called Eating Disorders

Neurodiversity Australia. However, no funding was obtained specifically for this narrative review.

Acknowledgements:

The authorship team would like to thank Phaedra Longhurst for her assistance reviewing the manuscript and providing feedback prior to its submission for publication.

REFERENCES

- 1. National Mental Health Commission. Our words matter: Guidelines for language use. Mindframe. 2023. https://mindframe.org.au/our-words-matter-guidelines-for-language-use
- 2. Bottema-Beutel K, Kapp SK, Lester JN, Sasson, NJ, Hand BN. Avoiding ableist language: Suggestions for autism researchers. Autism in adulthood. 2021;3. p. 18–29. https://doi.org/10.1089/aut.2020.0014
- 3. Monk R, Whitehouse AJO, Waddington H. The use of language in autism research. Trends in neurosciences. 2022;45. p. 791–793. https://doi.org/10.1016/j.tins.2022.08.009
- 4. Taboas A, Doepke K, Zimmerman C. Preferences for identity-first versus person-first language in a US sample of autism stakeholders. Autism. 2023;27. p. 565-570. https://doi.org/10.1177/13623613221130845
- 5. Kapp SK. Profound concerns about "profound autism": Dangers of severity scales and functioning labels for support needs. Educ Sci. 2023;13:106. https://doi.org/10.3390/educsci13020106
- 6. Natri HM, Abubakare O, Asasumasu K, Basargekar A, Beaud F, Botha M et al. Anti-Ableist language is fully compatible with high-quality autism research: Response to Singer et al. (2023). Autism Research. 2023;16. p. 673–676. https://doi.org/10.1002/aur.2928
- 7. Finesilver C, Leigh J, Brown N. Invisible disability, unacknowledged diversity. In Brown N, Leigh J, editors. Ableism in academia. London: UCL. 2020. p. 143-160.
- 8. Boyd V. 'Looking okay': exploring constructions of fluctuating or recurring impairments in UK higher education. Dissertation, university of Stirling. 2012.

- 9. Adams KL, Mandy W, Catmur C, Bird G. Potential mechanisms underlying the association between feeding and eating disorders and autism. Neurosci Biobehav Rev. 2024; 162:105717 doi:10.1016/j.neubiorev.2024.105717
- 10. Field SL, Fox JRE, Jones CRG, lliams MO.). "Work WITH us": A Delphi study about improving eating disorder treatment for autistic women with anorexia nervosa. J Eat Disord. 2023;11. https://doi.org/10.1186/s40337-023-00740-z
- 11. Apperly IA., Lee R, van der Kleij SW, Devine RT. A transdiagnostic approach to neurodiversity in a representative population sample: The N + 4 model. JCPP Adv. 2024. https://doi.org/10.1002/jcv2.12219
- 12. Fletcher-Watson S. Transdiagnostic research and the neurodiversity paradigm: Commentary on the transdiagnostic revolution in neurodevelopmental disorders by Astle et al. J Child Psychol Psychiatry. 2022;63. p. 418-420. https://doi.org/10.1111/jcpp.13589
- 13. Waldren LH, Leung FYN, Hargitai LD, Burgoyne AP., Liceralde, VRT, Livingston LA et al. Unpacking the overlap between Autism and ADHD in adults: A multi-method approach. Cortex. 2024;173. p. 120–137. https://doi.org/10.1016/j.cortex.2023.12.016
- 14. Hay P, Aouad P, Le A, Marks P, Maloney D; National Eating Disorder Research Consortium, et al. Epidemiology of eating disorders: population, prevalence, disease burden and quality of life informing public policy in Australia-a rapid review. J Eat Disord. 2023 Feb 15;11:1:23. doi: 10.1186/s40337-023-00738-7.
- 15. Pastore M, Indrio F, Bali D, Vural M, Giardino I, Pettoello-Mantovani M. Alarming increase of eating disorders in children and adolescents. J Pediatr. 2023;263.
- 16. Barakat S, McLean SA, Bryant E, Le A, Marks P, Touyz S et al. Risk factors for eating disorders: findings from a rapid review. J Eat Disord. 2023;11:8.

- 17. Solmi M, Monaco F, Højlund M, Monteleone AM, Trott M, Firth J et al. Outcomes in people with eating disorders: a transdiagnostic and disorder-specific systematic review, meta-analysis and multivariable meta-regression analysis. World Psychiatry. 2024;23. p. 124-138.
- 18. Miskovic-Wheatley J, Bryant E, Ong SH, Vatter S, Le, Touyz S, Maguire S. Eating disorder outcomes: findings from a rapid review of over a decade of research. J Eat Disord. 2023. p. 11:85. https://doi.org/10.1186/s40337-023-00801-3
- 19. Elwyn R. A lived experience response to the proposed diagnosis of terminal anorexia nervosa: Learning from iatrogenic harm, ambivalence and enduring hope. J Eat Disord. 2023;11. https://doi.org/10.1186/s40337-022-00729-0
- 20. Kenny TE, Lewis SP. More than an outcome: a person-centered, ecological framework for eating disorder recovery. J Eat Disord. 2023;11:45.
- 21. Wetzler S, Hackmann C, Peryer G, Clayman K, Friedman D, Saffran K et al. A framework to conceptualize personal recovery from eating disorders: A systematic review and qualitative meta-synthesis of perspectives from individuals with lived experience. Int J Eat Disord. 2020;53. p. 1188–1203. https://doi.org/10.1002/eat.23260
- 22. den Houting J. Neurodiversity: An insider's perspective. Autism. 2019;23. p. 271-273. https://doi.org/10.1177/1362361318820762
- 23. Legault M, Bourdon JN, Poirier P. From neurodiversity to neurodivergence: The role of epistemic and cognitive marginalization. Synthese. 2021;199. p. 12843-12868. https://doi.org/10.1007/s11229-021-03356-5
- 24. Giwa Onaiwu M. I, too, sing neurodiversity. Ought. 2020;2. https://doi.org/10.9707/2833-1508.1048

- 25. Leong C, Graichen R. Decentering Neuronormativity: A transactional analysis impasse theory perspective for understanding adhd masking and authentically honoring the da vinci archetype within. Transactional Analysis Journal. 2024;54. p. 91–106. https://doi.org/10.1080/03621537.2024.2286581
- 26. Bolton MJ. De-centering neuronormativity is an imperative in humanistic psychotherapy: Towards a neurodiversity-informed, person-centered approach. 2023. https://doi.org/10.31234/osf.io/q2t8h
- 27. Pellicano E, den Houting J. Annual research review: Shifting from 'normal science' to neurodiversity in autism science. J Child Psychol Psychiatry. 2022;63. p. 381–396. https://doi.org/10.1111/jcpp.13534
- 28. Chapman R. Neurodiversity and the social ecology of mental functions. Perspect Psychol Sci. 2021;16. p. 1360-1372. https://doi.org/10.1177/1745691620959833
- 29. Dwyer P. The neurodiversity approach (es): what are they and what do they mean for researchers? hum dev. 2022;66. p. 73–92. https://doi.org/10.1159/000523723
- 30. Lyons K. The neurodiversity paradigm and abolition of psychiatric incarceration. Columbia Law Review. 2023;123. p. 1993–2034. https://www.jstor.org/stable/27264955
- 31. Keski-Rahkonen A, Ruusunen A. Avoidant-restrictive food intake disorder and autism: Epidemiology, etiology, complications, treatment, and outcome. Curr Opin Psychiatry. 2023;36. p. 438-442. https://doi.org/10.1097/YCO.00000000000000896
- 32. Villa FM, Crippa A, Rosi E, Nobile M, Brambilla P, Delvecchio G. ADHD and eating disorders in childhood and adolescence: An updated minireview. J Affect Disord. 2023;321:265-271. doi:10.1016/j.jad.2022.10.016

- 33. Gal E, Hardal-Nasser R, Engel-Yeger B. The relationship between the severity of eating problems and intellectual developmental deficit level. Res Dev Disabil. 2011;32. p. 1464-1469. doi:10.1016/j.ridd.2010.12.003
- 34. Smith BL, Ludlow AK. An exploration of eating behaviours and caregiver mealtime actions of children with Tourette syndrome. Frontiers. 2022. https://doi.org/10.3389/fped.2022.933154
- 35. Bamigbade S-E, Rogers SL, Wills W, Ludlow AK. Mothers' accounts of mealtime and feeding challenges for children with tourette syndrome or persistent tic disorders. Front Psychiatry. 2022;13. https://doi.org/10.3389/fpsyt.2022.936796
- 36. Babb C, Brede J, Jones CRG, Serpell L, Mandy W, Fox J. A comparison of the eating disorder service experiences of autistic and non-autistic women in the UK. Eur Eat Disord Rev. 2022;30. p. 616–627. https://doi.org/10.1002/erv.2930
- 37. Svedlund NE, Norring C, Ginsberg Y, von Hausswolff-Juhlin Y. Symptoms of attention deficit hyperactivity disorder (ADHD) among adult eating disorder patients. BMC psychiatry. 2017;17:1-9.
- 38. Thomas E, Benjamin-Thomas TE, Sithambaram A, Shankar J, Chen SP. Participatory action research among people with serious mental illness: A scoping review. qual health res. 2024;34. p. 3–19. https://doi.org/10.1177/10497323231208111
- 39. Musić S, Elwyn R, Fountas G, Gnatt I, Jenkins Z M, Malcolm A et al. Valuing the voice of lived experience of eating disorders in the research process: benefits and considerations. Aus NZ J Psychiatry. 2022;56. p. 216-218.

- 40. Nimbley E, Maloney E, Duffy F. A call for autism-led research exploring definitions of recovery in Autistic individuals with an eating disorder. Int J Eat Disord. 2023;56. p. 1335–1340. https://doi.org/10.1002/eat.23953
- 41. Ambwani S, Coull E, Cardi V, Rowlands K, Treasure J. Every mistake is a treasure: Lessons learned from the TRIANGLE trial for anorexia nervosa. Int J Eat Disord. 2024;57. p. 1330-1336. https://doi.org/10.1002/eat.24121
- 42. Gillberg C. Are Autism and Anorexia Nervosa Related? Br J Psychiatry. 1983. p. 142:428. doi:10.1192/bjp.142.4.428b
- 43. Fisman S, Steele M, Short J, Byrne T, Lavallee C. Case Study: Anorexia Nervosa and Autistic Disorder in an Adolescent Girl. J Am Acad Child Adolesc Psychiatry. 1996. p. 35:937-940. doi:10.1097/00004583-199607000-00021
- 44. Gravestock S. Eating disorders in adults with intellectual disability. J Intellect Disabil Res. 2000;44:625–637. https://doi.org/10.1046/j.1365-2788.2000.00308.x
- 45. Surman CB, Randall ET, Biederman J. Association between attention-deficit/hyperactivity disorder and bulimia nervosa: analysis of 4 case-control studies. J Clin Psychiatry. 2006;67. p. 351–354. https://doi.org/10.4088/jcp.v67n0303
- 46. Lang J, Wylie G, Haig C, Gillberg C, Minnis H. Towards system redesign: An exploratory analysis of neurodivergent traits in a childhood population referred for autism assessment. PloS one. 2024;19. https://doi.org/10.1371/journal.pone.0296077
- 47. Cravedi E, Deniau E, Giannitelli M, Xavier J, Hartmann A, Cohen D. Tourette syndrome and other neurodevelopmental disorders: a comprehensive review. Child Adolesc. Psychiatry Ment. Health. 2017;11. p. 1-12.

- 48. Watts J. The epistemic injustice of borderline personality disorder. Brit J Psychiatry Int. 2024. p. 1–5. doi:10.1192/bji.2024.16
- 49. McQuaid GA, Strang JF, Jack A. Borderline personality as a factor in late, missed, and mis-diagnosis in autistic girls and women: A conceptual analysis. autism in adulthood. 2024. doi:10.1089/aut.2023.0034
- 50. Lai MC. Clinical reflections on the intersections of autism and personality development. Autism. 2022;26. p. 739–742. https://doi.org/10.1177/13623613221088073
- 51. Hartley S, Baker C, Birtwhistle M, Burgess JL, Chatburn E, Cobbaert L et al. Commentary: Bringing together lived experience, clinical and research expertise a commentary on the May 2022 debate (should CAMH professionals be diagnosing personality disorder in adolescence?). Child and Adolescent Mental Health. 2022;27. p. 246-249. https://doi.org/10.1111/camh.12586
- 52. Martin J. Why are females less likely to be diagnosed with ADHD in childhood than males? The Lancet Psychiatry. 2024;11. p. 303-310.
- 53. Murphy S, Flower RL, Jellett R. Women seeking an autism diagnosis in Australia: A qualitative exploration of factors that help and hinder. Autism. 2023;27. p. 808-821. https://doi.org/10.1177/13623613221117911
- 54. Rudra A. Why many women with autism and ADHD aren't diagnosed until adulthood and what to do if you think you're one of them. The Conversation. 2023. https://theconversation.com/why-many-women-with-autism-and-adhd-arent-diagnosed-until-adulthood-and-what-to-do-if-you-think-youre-one-of-them-179970

- 55. Kentrou V, Livingston LA, Grove R, Hoekstra RA, Begeer S. Perceived misdiagnosis of psychiatric conditions in autistic adults. eClinicalMedicine. 2024;71. https://doi.org/10.1016/j.eclinm.2024.102586
- 56. McCrossin R. Finding the true number of females with autistic spectrum disorder by estimating the biases in initial recognition and clinical diagnosis. Children. 2022;9:272. https://doi.org/10.3390/children9020272
- 57. Bailey B, Arciuli J. Indigenous Australians with autism: A scoping review. Autism. 2020;24. p. 1031–1046. https://doi.org/10.1177/1362361319894829
- 58. Bruno G, Chan TA., Zwaigenbaum L, Coombs E, Indigenous relations circle, Nicholas D. Indigenous autism in Canada: A scoping review. Journal of Autism and Developmental Disorders. 2023. https://doi.org/10.1007/s10803-023-06045-z
- 59. Garb HN. Race bias and gender bias in the diagnosis of psychological disorders. Clin Psych Rev. 2021;90:102087. https://doi.org/10.1016/j.cpr.2021.102087
- 60. Obeid R, Bisson JB, Cosenza A, Harrison AJ, James F, Saade S et al. Do implicit and explicit racial biases influence autism identification and stigma? An implicit association test study. J Autism Dev Disord. 2021;51. p. 106–128. https://doi.org/10.1007/s10803-020-04507-2
- 61. Mandell DS, Wiggins LD, Carpenter LA, Daniels J, DiGuiseppi C, Durkin MS et al. Racial/ethnic disparities in the identification of children with autism spectrum disorders. Am J Pub Health. 2009;99. p. 493–498. https://doi.org/10.2105/AJPH.2007.131243
- 62. Mandell DS, Ittenbach RF, Levy SE, Pinto-Martin JA. Disparities in diagnoses received prior to a diagnosis of autism spectrum disorder. J Autism Devel Disord. 2007;37. p. 1795–1802. https://doi.org/10.1007/s10803-006-0314-8

- 63. Ballentine KL. Understanding racial differences in diagnosing odd versus ADHD using critical race theory. Fam Soc. 2019;100. p. 282-292. https://doi.org/10.1177/1044389419842765
- 64. Fadus MC, Ginsburg KR, Sobowale K, Halliday-Boykins CA, Bryant BE, Gray KM et al. Unconscious bias and the diagnosis of disruptive behavior disorders and ADHD in African American and Hispanic Youth. Acad Psychiatry. 2020;44. p. 95–102. https://doi.org/10.1007/s40596-019-01127-6
- 65. Longhurst P, Todd J, Aspell JE, Swami V. Psychometric evaluation of a revised version of the body appreciation scale-2 for autistic adults (BAS-2A). Body Image. 2024;49. doi:10.1016/j.bodyim.2024.101706
- 66. Longhurst P, Nimbley E, Evans EH, Maclennan K, Gillespie-Smith K, Duffy F. Measuring eating disorders in autistic people: a proposal for future research. Manuscript under review. 2024.
- 67. Rong Y, Yang C, Jin Y, Wang Y. Prevalence of attention-deficit/hyperactivity disorder in individuals with autism spectrum disorder: A meta-analysis. Res autism spectrum disord. 2021;83. doi:10.1016/j.rasd.2021.101759
- 68. Huke, V, Turk, J, Saeidi, S, Kent, A, Morgan, JF. Autism spectrum disorders in eating disorder populations: A systematic review. European Eating Disorders Review; 2013;21. p. 345–351. doi:10.1002/erv.2244
- 69. Boltri M, Sapuppo W. Anorexia nervosa and autism spectrum disorder: a systematic review. Psychiatry Res. 2021;306:114271.

- 70. Koomar T, Thomas TR, Pottschmidt NR, Lutter M, Michaelson JJ. 2021). Estimating the prevalence and genetic risk mechanisms of ARFID in a large autism cohort. Frontiers. https://doi.org/10.3389/fpsyt.2021.668297
- 71. Clyde-Smith I, <u>Duffield</u>, C. Boy, 7, died of undiagnosed eating disorder that saw him living on just 'a few biscuits'. https://www.mirror.co.uk/news/uk-news/boy-7-died-undiagnosed-eating-31872750
- 72. Papini NM, Bulik CM, Chawner SJ, Micali N. Prevalence and recurrence of pica behaviors in early childhood: findings from the ALSPAC birth cohort. MedRxiv. 2023;23290951. https://doi.org/10.1101/2023.06.04.23290951
- 73. Fields VL, Soke GN, Reynolds A, Tian LH, Wiggins L, Maenner M et al. Pica, autism, and other disabilities. pediatrics. 2021;147. https://doi.org/10.1542/peds.2020-0462
- 74. Gesi C, Carmassi C, Luciano M, Bossini L, Ricca V, Fagiolini A et al. Autistic traits in patients with anorexia nervosa, bulimia nervosa or binge eating disorder: A Pilot Study. European Psychiatry. 2017;41. p. S100–S100. doi:10.1016/j.eurpsy.2017.01.310
- 75. Carpita B, Muti D, Cremone IM, Fagiolini A, Dell'Osso L. Eating disorders and autism spectrum: links and risks. CNS Spectrums. 2022;27. p. 272–280. doi:10.1017/S1092852920002011
- 76. Nazar BP, Bernardes C, Peachey G, Sergeant J, Mattos P, Treasure, J. The risk of eating disorders comorbid with attention-deficit/hyperactivity disorder: A systematic review and meta-analysis. Int J Eat Disord. 2016;49. p. 1045–1057. https://doi.org/10.1002/eat.22643

- 77. Adler LA, Spencer T, Faraone SV, Kessler RC, Howes MJ, Biederman J et al. Validity of pilot adult adhd self- report scale (ASRS) to rate adult ADHD symptoms. Ann Clin Psychiatry. 2006;18. p. 145–148. https://doi.org/10.3109/10401230600801077
- 78. Reinblatt SP, Mahone EM, Tanofsky-Kraff M, Lee-Winn AE, Yenokyan G, Leoutsakos JM. Pediatric loss of control eating syndrome: Association with attention-deficit/hyperactivity disorder and impulsivity. Int J Eat Disord. 2015;48. p. 580–588. https://doi.org/10.1002/eat.22404
- 79. Leib S, Gilon Mann T, Stein D, Vusiker I, Tokatly Latzer I, Ben-Ami, M et al. High prevalence of attention-deficit/hyperactivity disorder in adolescents with severe obesity seeking bariatric surgery. Acta Paediatr. Int. J. Paediatr. 2020;109. p. 581-586. doi:10.1111/apa.15039
- 80. Alfonsson S, Parling T, Ghaderi A. Screening of adult ADHD among patients presenting for bariatric surgery. Obes Surg. 2012;22. p. 918-926. doi:10.1007/s11695-011-0569-9
- 81. Cervantes PE, Matson JL. Comorbid symptomology in adults with autism spectrum disorder and intellectual disability. J Autism Devel Disord. 2015;45. p. 3961-3970. https://doi.org/10.1007/s10803-015-2553-z
- 82. Hove O. Prevalence of eating disorders in adults with mental retardation living in the community. Am J Ment Retard. 2004;109. p. 501–506. https://doi.org/10.1352/0895-8017(2004)109<501:POEDIA>2.0.CO;2
- 83. Manduchi B, Fainman GM, Walshe M. Interventions for feeding and swallowing disorders in adults with intellectual disability: A systematic review of the evidence.

 Dysphagia. 2020;35. p. 207-219. https://doi.org/10.1007/s00455-019-10038-5

- 84. Hirschtritt ME, Lee PC, Pauls DL, Dion Y, Grados MA, Illmann C. Lifetime prevalence, age of risk, and genetic relationships of comorbid psychiatric disorders in Tourette syndrome. JAMA Psychiatry. 2015;72. p. 325-333. https://doi.org/10.1001/jamapsychiatry.2014.2650
- 85. Drakes DH, Fawcett EJ, Rose JP, Carter-Major JC, Fawcett, JM. Comorbid obsessive-compulsive disorder in individuals with eating disorders: an epidemiological meta-analysis. J Psychiatr Res. 2021;141. p. 176-191.
- 86. Mandelli L, Draghetti S, Albert U, De Ronchi D, Atti AR. Rates of comorbid obsessive-compulsive disorder in eating disorders: a meta-analysis of the literature. J Affect Disord 2020;277. p. 927-939.
- 87. Lee EB, Barney JL, Twohig MP, Lensegrav-Benson T, Quakenbush B. Obsessive compulsive disorder and thought action fusion: Relationships with eating disorder outcomes. Eat Behav. 2020;37:101386.
- 88. Rodgers E, Marwaha S, Humpston C. Co-occurring psychotic and eating disorders in England: findings from the 2014 Adult Psychiatric Morbidity Survey. J Eat Disord. 2022;10:150.
- 89. Sankaranarayanan A, Johnson K, Mammen SJ, Wilding HE, Vasani D, Murali V et al. Disordered eating among people with schizophrenia spectrum disorders: A systematic review. Nutrients. 2021;13:3820.
- 90. Miotto P, Pollini B, Restaneo A, Favaretto G, Sisti D, Rocchi MB et al. Symptoms of psychosis in anorexia and bulimia nervosa. Psychiatry Res. 2010;175. p. 237-243.
- 91. Sarro S. Transient psychosis in anorexia nervosa: review and case report. Eating and Weight Disorders-Studies on Anorexia, Bulimia and Obesity. 2009;14. p. e139-e143.

- 92. Beisser SR, Gillespie CW. An investigation of eating disorders among gifted adolescents. Children and Teenagers. 2021;4. https://doi.org/10.22158/ct.v4n1p64
- 93. Bell AJ. The lived experience of an eating disorder among gifted female adolescents: a phenomenological study [Doctoral dissertation] University of British Columbia. 2004. https://open.library.ubc.ca/media/download/pdf/831/1.0099750/1
- 94. Garner D. Eating disorders in the gifted adolescent. Understanding the gifted adolescent. 1991:50-64.
- 95. Hughes CE. Giftedness and autism spectrum disorders. In Understanding twice-exceptional learners. Routledge. 2021. p. 217-243
- 96. Schilder CMT, van Elburg A A, Snellen WM, Sternheim LC, Hoek HW, Danner UN. Intellectual functioning of adolescent and adult patients with eating disorders. Int J Eat Disord. 2017;50. p. 481-489. https://doi.org/10.1002/eat.22594
- 97. Trail BA. Twice-exceptional learners: What they need in order to thrive. University of Northern Colorado. ProQuest Dissertations & Theses Global.
- 98. Blanz BJ, Detzner U, Lay B, Schmidt MH. The intellectual functioning of adolescents with anorexia nervosa and bulimia nervosa. Eur Child Adolesc Psychiatry. 1997;6:129-135. https://doi.org/10.1007/BF00538984
- 99. Murray D, Milton D, Green J, Bervoets J. The human spectrum: A phenomenological enquiry within neurodiversity. Psychopathology. 2023;56:220-230. https://doi.org/10.1159/000526213
- 100. Green J, Shaughnessy N. Autistic phenomenology: past, present, and potential future. Front Psychol. 2023;14:1287209. https://doi.org/10.3389/fpsyg.2023.1287209

- 101. Cobbaert L, Rose A. Eating disorders and neurodivergence: A stepped care approach.2023
- 102. Longhurst P. Body image and autism: A scoping review. Research in Autism Spectrum Disorders. 2023;105: 102170. https://doi.org/10.1016/j.rasd.2023.102170
- 103. Longhurst P, Burnette CB. Challenges and opportunities for conceptualizing intuitive eating in autistic people. Int J Eat Disord. 2023;56. p. 2189–2199.

https://doi.org/10.1002/eat.24057

- 104. Dellapiazza F, Michelon C, Vernhet C, Muratori F, Blanc N, Picot MC, et al. Sensory processing related to attention in children with ASD, ADHD, or typical development: Results from the ELENA cohort. Eur Child Adolesc Psychiatry. 2021;30. p. 283-291. https://doi.org/10.1007/s00787-020-01516-5
- 105. Gere DR, Capps SC, Mitchell DW, Grubbs E. Sensory sensitivities of gifted children. Am J Occup Ther. 2009;63. p. 288-295. https://doi.org/10.5014/ajot.63.3.288
- 106. Houghton DC, Capriotti MR, Conelea CA, Woods DW. Sensory phenomena in Tourette syndrome: Their Role in Symptom Formation and Treatment. Curr Dev Disord Rep. 2014;1. p. 245–251. https://doi.org/10.1007/s40474-014-0026-2
- 107. Tran HT, Li YC, Lin HY, Lee SD, Wang PJ. Sensory processing impairments in children with developmental coordination disorder. Children. 2022;9:10:1443. https://doi.org/10.3390/children9101443
- 108. Viana AR, Razuk M, de Freitas PB, Barela JA. Sensorimotor integration in dyslexic children under different sensory stimulations. PloS one. 2013;8: e72719. https://doi.org/10.1371/journal.pone.0072719

- 109. Cervin M. Sensory processing difficulties in children and adolescents with obsessive-compulsive and anxiety disorders. Res Child Adolesc Psychopathol. 2023;51. p. 223–232. https://doi.org/10.1007/s10802-022-00962-w
- 110. Ramsay IS, Schallmo M-P, Biagianti B, Fisher M, Vinogradov S, Sponheim SR. Deficits in auditory and visual sensory discrimination reflect a genetic liability for psychosis and predict disruptions in global cognitive functioning. Front Psychiatry. 2020;11. https://doi.org/10.3389/fpsyt.2020.00638
- 111. Burke E, O'Dwyer M, Maes-Festen D, Oppewal A. Chronic health among those with an intellectual disability. In F. Sheerin & C. Doyle (Eds.), Intellectual disabilities: Health and social care across the lifespan. Springer; 2023. https://doi.org/10.1007/978-3-031-27496-1_9
- 112. Sibeoni J, Massoutier L, Valette M, Manolios E, Verneuil L, Speranza M, et al. The sensory experiences of autistic people: A metasynthesis. Autism. 2022;26:1032-1045.
- 113. Cobbaert L, Hay P, Mitchell PB, Roza SJ, Perkes I. Sensory processing across eating disorders: A systematic review and meta-analysis of self-report inventories. Int J Eat Disord. 2024. https://doi.org/10.1002/eat.24184
- 114. Navas-León S, Tajadura-Jiménez A, Motrico E, Morales L, Borda-Mas M, Almeda N, Sánchez-Martín M. Understanding and treating body image disturbances in eating disorders through body illusion interventions: A scoping review protocol. Syst Rev. 2024;13:65. https://doi.org/10.1186/s13643-024-02458-8
- 115. Saure E, Lepistö-Paisley T, Raevuori A, Laasonen M. Atypical sensory processing is associated with lower body mass index and increased eating disturbance in individuals with anorexia nervosa. Front Psychiatry. 2022;13:850594.

- 116. Nimbley E, Golds L, Sharpe H, Gillespie-Smith K, Duffy F. Sensory processing and eating behaviours in autism: A systematic review. Eur Eat Disord Rev. 2022;30. p. 538–559. https://doi.org/10.1002/erv.2920
- 117. Nisticò V, Faggioli R, Tedesco R, Giordano B, Priori A, Gambini O, et al. Brief report: sensory sensitivity is associated with disturbed eating in adults with autism spectrum disorders without intellectual disabilities. J Autism Dev Disord. 2023;53. p. 3295-3300.
- 118. Elsayed HE, Thompson KL, Conklin JL, Watson, LR. Systematic review of the relation between feeding problems and sensory processing in children with autism spectrum disorder. Am J Speech Lang Pathol. 2022;31. p. 2875-2899.
- 119. Wolff M, Morceau S, Folkard R, Martin-Cortecero J, Groh A. A thalamic bridge from sensory perception to cognition. Neurosci Biobehav Rev. 2021;120. p. 222-235. doi:10.1016/j.neubiorev.2020.11.013
- 120. Wolff M, Vann SD. The cognitive thalamus as a gateway to mental representations. J Neurosci. 2019;39. p. 3–14. https://doi.org/10.1523/JNEUROSCI.0479-18.2018
- 121. Saalmann YB, Kastner S. The cognitive thalamus. Front Syst Neurosci. 2015;9:39.
- 122. Cabrera-Álvarez J, Doorn N, Maestú F, Susi G. Modeling the role of the thalamus in resting-state functional connectivity: Nature or structure. PLoS Comput Biol. 2023;19:8. https://doi.org/10.1371/journal.pcbi.1011007
- 123. Fama R, Sullivan EV. Thalamic structures and associated cognitive functions: Relations with age and aging. Neuroscience and biobehavioral reviews. 2015;54. p. 29–37. https://doi.org/10.1016/j.neubiorev.2015.03.008

- 124. Wen X, Li W, Liu Y, et al. Exploring communication between the thalamus and cognitive control-related functional networks in the cerebral cortex. Cogn Affect Behav Neurosci. 2021;2:13. p. 656-677. doi:10.3758/s13415-021-00892-y.
- 125. Ouhaz Z, Fleming H, Mitchell AS. Cognitive functions and neurodevelopmental disorders involving the prefrontal cortex and mediodorsal thalamus. Front Neurosci. 2018 Feb 6;12:33. doi: 10.3389/fnins.2018.00033.
- 126. Barson JR, Mack NR, Gao WJ. The paraventricular nucleus of the thalamus is an important node in the emotional processing network. Front Behav Neurosci. 2020;29:14. p. 598469. doi: 10.3389/fnbeh.2020.598469.
- 127. Hwang WJ, Kwak YB, Cho KIK, Lee TY, Oh H, Ha M, Kim M, Kwon JS. Thalamic connectivity system across psychiatric disorders: current status and clinical implications. Biol Psychiatry Glob Open Sci. 2021;2(4). p. 332-340.

https://doi.org/10.1016/j.bpsgos.2021.09.008

- 128. Yang B, Jia Y, Zheng W, Wang L, Qi Q, Qin W, Li X, Chen X, Lu J, Li H, Zhang Q, Chen N. Structural changes in the thalamus and its subregions in regulating different symptoms of posttraumatic stress disorder. Psychiatry Res Neuroimaging. 2023;335:111706. doi: 10.1016/j.pscychresns.2023.111706.
- 129. Kirouac GJ. The paraventricular nucleus of the thalamus as an integrating and relay node in the brain anxiety network. Front Behav Neurosci. 2021;24;15:627633. doi: 10.3389/fnbeh.2021.627633.
- 130. Wang J, Wu G, Wang M, Li W, Wang Y, Ren X, et al. Exploring the thalamus: a crucial hub for brain function and communication in patients with bulimia nervosa. J Eat Disord. 2023;11:207. doi:10.1186/s40337-023-00933-6.

- 131. Petrovich GD. The function of paraventricular thalamic circuitry in adaptive control of feeding behavior. Front Behav Neurosci. 2021 Apr 26;15:671096. doi: 10.3389/fnbeh.2021.671096.
- 132. Millan EZ, Ong Z, McNally GP. Chapter 5 Paraventricular thalamus: Gateway to feeding, appetitive motivation, and drug addiction. In: Calvey T, Daniels WMU, editors. Progress in Brain Research. Elsevier. 2017. p. 113-137. doi: 10.1016/bs.pbr.2017.07.006.
- 133. Gavnik AA. A sociological examination of disordered eating and body satisfaction among transgender and nonbinary individuals. [Master's thesis]. University of Delaware; 2023.
- 134. Park-Cardoso J, da Silva APS. Preference to Eat Alone: Autistic Adults' Desire for Freedom of Choice for a Peaceful Space. Autism in Adulthood. 2021;3. p. 257-265. doi:10.1089/aut.2020.0066
- 135. Kluckow H, Telfer J, Abraham S. Should we screen for misophonia in patients with eating disorders? A report of three cases. Int J Eat Disord. 2014;47. p. 558-561. https://doi.org/10.1002/eat.22245
- 136. Naraindas AM, Moreno M, Cooney SM. Beyond Gender: Interoceptive Sensibility as a Key Predictor of Body Image Disturbances. Behav Sci. 2023;14:25. https://doi.org/10.3390/bs14010025
- 137. Todd J, Aspell JE, Barron D, Swami V. Multiple dimensions of interoceptive awareness are associated with facets of body image in British adults. Body Image. 2019;29. p. 6-16. doi:10.1016/j.bodyim.2019.02.003

- 138. Lyvers M, Kelahroodi M, Udodzik E, Stapleton P, Thorberg FA. Alexithymia and binge eating: Maladaptive emotion regulation strategy or deficient interoception? Appetite. 2022;175:106073. doi:10.1016/j.appet.2022.106073
- 139. Wallis DJ, Ridout N. Direct and indirect effects of alexithymia on disordered eating in a non-clinical female sample: Determining the role of negative affect. Front Psychiatry. 2022 Nov 24;13:994024. doi: 10.3389/fpsyt.2022.994024.
- 140. Vuillier L, Carter Z, Teixeira AR, Moseley RL. Alexithymia may explain the relationship between autistic traits and eating disorder psychopathology. Mol Autism. 2020;11. p. 1-19.
- 141. Kinnaird E, Norton C, Pimblett C, Stewart C, Tchanturia K. Eating as an autistic adult: An exploratory qualitative study. PloS one. 2019;14: e0221937. https://doi.org/10.1371/journal.pone.0221937
- 142. Park-Cardoso J, da Silva APS. Insistence on sameness for food space appropriation: An exploratory study on Brazilians with autism (self-)diagnosis in adulthood. Autism. 2023;27. p. 938-951. https://doi.org/10.1177/13623613221121417
- 143. Jamal W, Cardinaux A, Haskins AJ, Kjelgaard M, Sinha P. Reduced sensory habituation in autism and its correlation with behavioral measures. J Autism Dev Disord. 2021;51. p. 3153–3164. https://doi.org/10.1007/s10803-020-04780-1
- 144. Cary E, Pacheco D, Kaplan-Kahn E, McKernan E, Matsuba E, Prieve B, et al. Brain signatures of early and late neural measures of auditory habituation and discrimination in autism and their relationship to autistic traits and sensory overresponsivity. J Autism Dev Disord. 2024;54. p. 1344–1360. https://doi.org/10.1007/s10803-022-05866-8

- 145. Kerns CM, Lankenau S, Shattuck PT, Robins DL, Newschaffer CJ, Berkowitz SJ.
 Exploring potential sources of childhood trauma: A qualitative study with autistic adults and caregivers. Autism. 2022;26. p. 1987–1998. https://doi.org/10.1177/13623613211070637
 146. Cristofori I, Cohen-Zimerman S, Grafman J. Chapter 11 Executive functions.
- Handbook of Clinical Neurology. 2019;163. p. 197-219. doi:10.1016/B978-0-12-804281-6.00011-2
- 147. Mareva S, Lead Investigators, Astle D, Baker K, Gathercole S, Holmes J, et al. Mapping neurodevelopmental diversity in executive function. Cortex. 2024;172. p. 204-221. doi:10.1016/j.cortex.2023.11.021
- 148. Diaz-Marsa M, Pemau A, de la Torre-Luque A, Vaz-Leal F, Rojo-Moreno L, Beato-Fernandez L, et al. Executive dysfunction in eating disorders: Relationship with clinical features. Prog. Neuropsychopharmacol. Biol. Psychiatry. 2023;120. doi:10.1016/j.pnpbp.2022.110649
- 149. Hirst RB, Beard CL, Colby KA, Quittner Z, Mills BM., Lavender JM. Anorexia nervosa and bulimia nervosa: A meta-analysis of executive functioning. Neurosci Biobehav Rev. 2017;83. p. 678–690. https://doi.org/10.1016/j.neubiorev.2017.08.011
- 150. Vicente A, Michel C, Petrolini V. Literalism in autistic people: A predictive processing proposal. Review of Philosophy and Psychology. 2023. https://doi.org/10.1007/s13164-023-00704-x
- 151. Brede J, Babb C, Jones C, Elliott M, Zanker C, Tchanturia K, et al. "For me, the anorexia is just a symptom, and the cause is the autism": Investigating restrictive eating disorders in autistic women. J Autism Dev Disord. 2020;50. p. 4280–4296. https://doi.org/10.1007/s10803-020-04479-3

- 152. Dell'Osso L, Nardi B, Benedetti F, Cremone IM, Casagrande D, Massimetti G, et al. Orthorexia and autism spectrum in University workers: relationship with gender, body mass index and dietary habits. Eat Weight Disord. 2022;27:8. p. 3713-3723. doi: 10.1007/s40519-022-01514-3.
- 153. Bouzy J, Brunelle J, Cohen D, Condat A. Transidentities and autism spectrum disorder: A systematic review. Psychiatry Res. 2023;323:115176.
- 154. Cheung AS, Ooi O, Leemaqz S, Cundill P, Silberstein N, Bretherton I, et al. Sociodemographic and clinical characteristics of transgender adults in Australia. Transgender health. 2018;3. p. 229-238.
- 155. De Vries AL, Roehle R, Marshall L, Frisén L, Van De Grift TC, Kreukels BP, et al. Mental health of a large group of adults with disorders of sex development in six European countries. Psychosom Med. 2019;81. p. 629-640.
- 156. George R, Stokes MA. Sexual orientation in autism spectrum disorder. Autism Res. 2018;11:133–141. https://doi.org/10.1002/aur.1892
- 157. Tollit MA, Maloof T, Hoq M, Haebich K, Pace CC, Rodriguez ZM., et al. A comparison of gender diversity in transgender young people with and without autistic traits from the Trans 20 cohort study. The Lancet Regional Health–Western Pacific. 2024:47.
- 158. Voltaire S, Steinberg H, Garfield T, Chvasta K, Ardeleanu K, Brown M, et al. Inextricably tied: Nonbinary autistic individuals' views on how their gender identity and autism are connected. Autism. 2024:13623613241257600.

https://doi.org/10.1177/13623613241257600

- 159. Weir E, Allison C, Baron-Cohen S. The sexual health, orientation, and activity of autistic adolescents and adults. Autism Res. 2021;14. p. 2342–2354. https://doi.org/10.1002/aur.2604
- 160. Cusack CE, Silverstein S, Askew AJ, Simone M, Galupo MP, Levinson CA. Eating disorders among queer and trans individuals: Implications for conceptualization, assessment, and treatment. Bulletin of the Menninger Clinic. 2024;88. p. 128-147.
- 161. Straus, P, Cook A, Watson V, Winter S, Whitehouse A, Albrecht N, et al. Mental health difficulties among trans and gender diverse young people with an autism spectrum disorder (ASD): Findings from Trans Pathways. J Psychiatr Res. 2021;137. p. 360-367.
- 162. Coelho JS, Suen J, Clark BA, Marshall SK, Geller J, Lam PY. Eating disorder diagnoses and symptom presentation in transgender youth: a scoping review. Curr Psychiatry Rep. 2019;21. p. 1-10.
- 163. Heiden-Rootes K, Linsenmeyer W, Levine S, Oliveras M, Joseph, M. A scoping review of the research literature on eating and body image for transgender and nonbinary adults. J Eat Disord. 2023;11:111.
- 164. Rasmussen SM, Dalgaard MK, Roloff M, Pinholt M, Skrubbeltrang C, Clausen L, et al. Eating disorder symptomatology among transgender individuals: a systematic review and meta-analysis. J Eat Disord. 2023;11:84.
- 165. Alvå H. "The greater dysphoria I've had, the greater dysmorphia-they work together"-An interview study on trans and nonbinary experiences of seeking and participating in Swedish eating disorder treatment. 2023.
- 166. Kester BL. Internalized transphobia and the development of disordered eating behaviors in gender diverse adults [Doctoral dissertation] Antioch University; 2023.

- 167. Bialy K. The Impact of gender minority stress on eating pathology and body checking behaviors in transgender and non-binary adults. [Dissertation] Philadelphia College of Osteopathic Medicine. 2023.
- 168. Biefield SD. Body-based harassment and eating disorder symptomology in cisgender, transgender, and gender nonconforming individuals. [Doctoral dissertation] University of Kentucky. 2023.
- 169. Soulliard ZA, Le TP, Yamasaki V. Pride and body image among transgender and nonbinary adults: the mediating role of body appreciation between gender identity pride and intuitive eating. Eating Disorders. 2024. p. 1-18.
- 170. McGregor K, Williams CR, Botta A, Mandel F, Gentile J. Providing essential gender-affirming telehealth services to transgender youth during COVID-19: A service review. J Telemed Telecare. 2023;29:147-152.
- 171. Velez, BL, Breslow AS, Brewster ME, Cox Jr R, Foster AB. Building a pantheoretical model of dehumanization with transgender men: Integrating objectification and minority stress theories. J Couns Psychol. 2016;63:497.
- 172. Cooper K, Mandy W, Butler C, Russell A. The lived experience of gender dysphoria in autistic adults: An interpretative phenomenological analysis. Autism. 2022;26. p. 963–974. https://doi.org/10.1177/13623613211039113
- 173. Cooper K, Mandy W, Butler C, Russell A. Phenomenology of gender dysphoria in autism: a multiperspective qualitative analysis. J Child Psychol Psychiatr. 2023;64:265-276. https://doi.org/10.1111/jcpp.13691

- 174. Elwyn R, Williams M, Smith E, Smith S. Two identical twin pairs discordant for longstanding anorexia nervosa and OSFED: Lived experience accounts of eating disorder and recovery processes. [Manuscript submitted for publication]. 2024.
- 175. Lundberg T, Roen K, Kraft C, Hegarty P. How young people talk about their variations in sex characteristics: making the topic of intersex talkable via sex education. Sex Educ. 2021;21:5. p. 552-567. doi:10.1080/14681811.2021.1911796.
- 176. Reis SM, Renzulli JS. Intellectual giftedness. In R. J. Sternberg (Ed.), The Cambridge handbook of intelligence (2nd ed.). Cambridge University Press; 2020: p. 291–316 https://doi.org/10.1017/9781108770422.014
- 177. Andronaco JA, Shute R, McLachlan A. Exploring asynchrony as a theoretical framework for understanding giftedness: A case of cognitive dissonance? Roeper Rev. 2014;36. p. 264-272.
- 178. Petersson S, Johnsson P, Perseius KI. A Sisyphean task: experiences of perfectionism in patients with eating disorders. J Eat Disord. 2017;5. p. 1-11.
- 179. Rice KG, Ray ME. Perfectionism and the gifted. In Pfeiffer SI, Shaunessy-Dedrick E, Foley-Nicpon M, editors, APA handbook of giftedness and talent (pp.). American Psychological Association; 2018. p. 645–658. https://doi.org/10.1037/0000038-042
- 180. Eli K. Striving for liminality: Eating disorders and social suffering. Transcult Psychiatry. 2018;55. p. 475-494.
- 181. Krafchek J, Kronborg L. The social coping of academically high-achieving females before the onset of disordered eating. Gifted Talented Int. 2020;35. p. 86–99. https://doi.org/10.1080/15332276.2020.1818329

- 182. Krafchek J, Kronborg L. Academic emotions experienced by academically high-achieving females who developed disordered eating. Roeper Rev. 2019;41. p. 258–272. https://doi.org/10.1080/02783193.2019.1661053
- 183. Peterson JS. Giftedness, trauma, and development: A qualitative, longitudinal case study. J Educ Gifted. 2014;37. p. 295-318.
- 184. Stackpole R, Greene D, Bills E, Egan SJ.). The association between eating disorders and perfectionism in adults: A systematic review and meta-analysis. Eat Behav. 2023;50. doi:101769. 10.1016/j.eatbeh.2023.101769
- 185. Townend G, Pendergast D, Garvis S. Academic self-concept in twice-exceptional students: What the literature tells us. TalentEd. 2014;28. p. 75–89. https://search.informit.org/doi/10.3316/informit.148086039194781
- 186. Amend ER, Peters DB. The importance of accurate assssment of gifted students: Issues with misdiagnosis, missed diagnoses, and twice-exceptionality. In handbook for counselors serving students with gifts and talents. routledge. 2021. p. 713-731
- 187. Desvaux T, Danna J, Velay JL, Frey A. From gifted to high potential and twice exceptional: A state-of-the-art meta-review. Appl. Neuropsychol. Child. 2023;13. p. 165–179. https://doi.org/10.1080/21622965.2023.2252950
- 188. Borsotti V, Begel A, Bjørn P. Neurodiversity and the accessible university: Exploring organizational barriers, access labor and opportunities for change. Proceedings of the ACM on Human-Computer Interaction, 8(CSCW1). 2024. p. 1–27. https://doi.org/10.1145/3641011
- 189. Arnold SR, Bruce G, Weise J, Mills CJ, Trollor JN, Coxon K. Barriers to healthcare for Australian autistic adults. Autism. 2023;28. p. 301-315.

https://doi.org/10.1177/13623613231168444

- 190. Tan DW, Rabuka M, Haar T, Pellicano E. 'It's a symbolic violence': Autistic people's experiences of discrimination at universities in Australia. Autism. 2024;28. p. 1344-1356. https://doi.org/10.1177/13623613231219744
- 191. Morton HE, Gillis JM, Zale EL, Brimhall KC, Romanczyk RG. Development and validation of the assessment of bullying experiences questionnaire for neurodivergent youth.

 J Autism Dev Disord. 2022. p. 1-14.
- 192. Mellifont D. Ableist ivory towers: a narrative review informing about the lived experiences of neurodivergent staff in contemporary higher education. Disability & Society. 2021;38. p. 865–886. https://doi.org/10.1080/09687599.2021.1965547
- 193. Praslova LN. Autism doesn't hold people back at work. discrimination does. Harv Bus Rev. 2021. https://hbr.org/2021/12/autism-doesnt-hold-people-back-at-work-discrimination-does
- 194. Moeller M, Ott DL., Russo, E. Neurodiversity can be a workplace strength, if we make room for it. The Conversation. 2021. https://theconversation.com/neurodiversity-can-be-a-workplace-strength-if-we-make-room-for-it-164859
- 195. Weitlauf AS, Miceli A, Vehorn A, Dada Y, Pinnock T, Harris JW, et al. Screening, diagnosis, and intervention for Autism: Experiences of black and multiracial families seeking care. J Autism Dev Disord. 2023. https://doi.org/10.1007/s10803-022-05861-z.
- 196. Smethurst, L., Thompson, A. R., & Freeth, M. (2024). "I've absolutely reached rock bottom and have no energy": The lived experience of unemployed and underemployed autistic adults. Autism in Adulthood.

- 197. Otiniano Verissimo AD, Henley N, Gee G, Davis CM. Homelessness and discrimination among US adults: the role of intersectionality. Journal of Social Distress and the Homeless. 2021;32:1. p. 1-15. DOI:10.1080/10530789.2021.1935650
- 198. Karpur A, Vasudevan V, Frazier TW, Shih AJ. Food insecurity in households of children with ASD in COVID-19 pandemic: A comparative analysis with the Household Pulse Survey data using stabilized inverse probability treatment weights. Disabil Health J. 2022;15:3. doi:10.1016/j.dhjo.2022.101323.
- 199. Moore R, Dada S, Emmambux MN, Samuels A. Food and nutrition security in persons with disabilities. A scoping review. Global Food Security. 2021;31:100581.
- 200. Kirkbride JB, Anglin DM, Colman I, Dykxhoorn J, Jones PB, Patalay P, et al. The social determinants of mental health and disorder: evidence, prevention and recommendations. World Psychiatry. 2024;23:1. p. 58-90. doi:10.1002/wps.21160.
- 201. Mason TB, Mozdzierz P, Wang S, Smith KE. Discrimination and eating disorder psychopathology: A meta-analysis. Behav Ther. 2021;52:2. p. 406-417. doi:10.1016/j.beth.2020.05.003.
- 202. Cleary M, West S, Kornhaber R, Hungerford C. Autism, discrimination and masking: Disrupting a recipe for trauma. Issues Ment Health Nurs. 2023;44:9. p. 799-808.
- 203. Davies J, Heasman B, Livesey A, Walker A, Pellicano E, Remington A. Access to employment: A comparison of autistic, neurodivergent and neurotypical adults' experiences of hiring processes in the United Kingdom. Autism. 2023;27:6. p. 1746-1763.
- 204. Gallagher A. Homelessness amongst autistic people and people with ADHD: a systematic review of the prevalence rates and risk factors and a qualitative exploration of individual experiences [Doctoral thesis]. University of Glasgow, School of Health and

Wellbeing, College of Medical, Veterinary and Life Sciences; 2023. Available from: https://theses.gla.ac.uk/83816/5/2023GallagherDClinPsy.pdf

- 205. Yalım E, Ünsal C, Gündoğmuş İ. The relationship between occupational functionality and metacognition in patients with obsessive-compulsive disorder. Cureus. 2024;16:1. doi:10.7759/cureus.51738.
- 206. Brown M, McCann E. Homelessness and people with intellectual disabilities: A systematic review of the international research evidence. J Appl Res Intellect Disabil. 2021;34:2. p. 390-401. doi:10.1111/jar.12815.
- 207. Hallward L, Nagata JM, Testa A, Jackson DB, Ganson KT. Associations between gender identity, eating disorder psychopathology, and food insecurity among Canadian adolescents and young adults during the COVID-19 pandemic. Eat Behav. 2023;49:101723.
- 208. Hazzard VM, Loth KA, Hooper L, Becker CB. Food insecurity and eating disorders: a review of emerging evidence. Curr Psychiatry Rep. 2020;22:12:74. doi:10.1007/s11920-020-01200-0.
- 209. Hazzard VM, Hooper L, Larson N, Loth KA, Wall MM, Neumark-Sztainer D. Associations between severe food insecurity and disordered eating behaviors from adolescence to young adulthood: Findings from a 10-year longitudinal study. Prev Med. 2022;154:106895. doi:10.1016/j.ypmed.2021.106895.
- 210. Han E, Scior K, Avramides K, Crane L. A systematic review on autistic people's experiences of stigma and coping strategies. Autism Res. 2022 Jan;15:1. p. 12-26. doi: 10.1002/aur.2652.

- 211. Masuch TV, Bea M, Alm B, et al. Internalized stigma, anticipated discrimination and perceived public stigma in adults with ADHD. ADHD Atten Def Hyp Disord. 2019;11. p. 212-220. doi:10.1007/s12402-018-0274-9.
- 212. Turnock A, Langley K, Jones CRG. Understanding stigma in autism: A narrative review and theoretical model. Autism Adulthood. 2022;4:1. p. 76-91. doi:10.1089/aut.2021.0005.
- 213. Hume R, Burgess H. "I'm Human After All": autism, trauma, and affective empathy. Autism Adulthood. 2021;3:3. p. 221-229.
- 214. Kimber L, Verrier D, Connolly S. Autistic people's experience of empathy and the autistic empathy deficit narrative. Autism Adulthood. 2023. doi:10.1089/aut.2023.0001.
- Milton D, Gurbuz E, López B. The 'double empathy problem': Ten years on. Autism.
 2022;268. p. 1901-1903. doi:10.1177/13623613221129123.
- 216. Gernsbacher MA, Yergeau M. Empirical failures of the claim that autistic people lack a theory of mind. Arch Sci Psychol. 2019;7. p. 102-118. https://doi.org/10.1037/arc0000067
- 217. Flanigan LK. I do not have stigma towards people with ADHD (but I do think they're lazy): Using education and experience to reduce negative attitudes towards ADHD (Doctoral thesis, University of Calgary, Calgary, Canada). (2021). Retrieved from https://prism.ucalgary.ca. http://hdl.handle.net/1880/113101
- 218. Shaw H. "I Don't Suffer from ADHD, I Suffer from Other People": ADHD, Stigma, and Academic Life. ADHD (Doctoral thesis, Dalhousie University, Canada). (2024).

 Retrieved from https://dalspace.library.dal.ca/handle/10222/80523
- 219. Radulski B. What are "masking" and "camouflaging" in the context of autism and ADHD? The Conversation. Published January 3, 2023. Available from:

https://theconversation.com/what-are-masking-and-camouflaging-in-the-context-of-autism-and-adhd-193446

- 220. Howe SJ, Hull L, Sedgewick F, Hannon B, McMorris CA. Understanding camouflaging and identity in autistic children and adolescents using photo-elicitation. Res Autism Spectr Disord. 2023;108:102232. doi:10.1016/j.rasd.2023.102232
- 221. Biberdzic M, Tang J, Tan J. Beyond difficulties in self-regulation: the role of identity integration and personality functioning in young women with disordered eating behaviours. J Eat Disord. 2021;9:93. https://doi.org/10.1186/s40337-021-00398-5
- 222. Evans JA, Krumrei-Mancuso E, Rouse SV. What You Are Hiding Could Be Hurting You: Autistic Masking in Relation to Mental Health, Interpersonal Trauma, Authenticity, and Self-Esteem. Autism Adulthood. 2023. doi:10.1089/aut.2022.0115
- 223. Cassidy SA, Gould K, Townsend E, Pelton M, Robertson AE, Rodgers J. Is camouflaging autistic traits associated with suicidal thoughts and behaviours? Expanding the interpersonal psychological theory of suicide in an undergraduate student sample. J Autism Dev Disord. 2020;50:10. p. 3638-3648. doi: 10.1007/s10803-019-04323-3.
- 224. Radulski EM. Conceptualising autistic masking, camouflaging, and neurotypical privilege: Towards a minority group model of neurodiversity. Human Development. 2022;66:2. p. 113-127. https://doi.org/10.1159/000524122
- 225. South M, Costa AP, McMorris C. Death by suicide among people with autism: Beyond zebrafish. JAMA Netw Open. 2021;4:1. doi:10.1001/jamanetworkopen.2020.34018
- 226. Bradley S, Moore F, Duffy F, Clark L, Suratwala T, Knightsmith P, Gillespie-Smith K. Camouflaging, not sensory processing or autistic identity, predicts eating disorder symptoms in autistic adults. Autism. 2024. doi:10.1177/13623613241245749

- 227. Douglas S, Sedgewick F. Experiences of interpersonal victimization and abuse among autistic people. Autism. 2023. doi:10.1177/13623613231205630
- 228. Pearson A, Rose K, Rees J. 'I felt like I deserved it because I was autistic': Understanding the impact of interpersonal victimisation in the lives of autistic people. Autism. 2023;27:2. p. 500-511. https://doi.org/10.1177/13623613221104546
- 229. Wymbs BT, Gidycz CA. Examining link between childhood ADHD and sexual assault victimization. J Atten Disord. 2021;25:11. p. 1612-1622. https://doi.org/10.1177/1087054720923750
- 230. Snyder JA. Examining the link between assault victimization and ADHD among college men. Crim Justice Stud. 2018;32:1. p. 16-31. https://doi.org/10.1080/1478601X.2018.1555159
- 231. Snyder JA. The link between ADHD and the risk of sexual victimization among college women: Expanding the lifestyles/routine activities framework. Violence Against Women. 2015;21:11. p. 1364-1384. https://doi.org/10.1177/1077801215593647
- 232. Cazalis F, Reyes E, Leduc S, Gourion D. Evidence that nine autistic women out of ten have been victims of sexual violence. Front Behav Neurosci. 2022;16:852203. https://doi.org/10.3389/fnbeh.2022.852203
- 233. Christian A. Majority of women with intellectual disability have experienced sexual abuse. Disability Support Guide. Published October 15, 2021. Available from:

 https://www.disability-have-experienced-sexual-abuse-hears-disability-royal-commission
- 234. Langevin R, Marshall C, Wallace A, Gagné M-E, Kingsland E, Temcheff C.
 Disentangling the associations between attention deficit hyperactivity disorder and child

sexual abuse: A systematic review. Trauma Violence Abuse. 2023;24:2. p. 369-389. https://doi.org/10.1177/15248380211030234

- 235. Mitchell KS, Scioli ER, Galovski T, Belfer PL, Cooper Z. Posttraumatic stress disorder and eating disorders: maintaining mechanisms and treatment targets. Eat Disord. 2021;29:3. p. 292-306. https://doi.org/10.1080/10640266.2020.
- 236. Boswell RG, Launius KN, Lydecker JA. Multiple marginalization, discrimination, and disordered eating among youth aged 10–11. Int J Eat Disord. 2024.
- 237. Nelson JD, Martin LN, Izquierdo A, Kornienko O, Cuellar AE, Cheskin LJ, Fischer S. The role of discrimination and adverse childhood experiences in disordered eating. J Eat Disord. 2023;11:1:29. https://doi.org/10.1186/s40337-023-00753-8
- 238. Lobregt-van Buuren E, Hoekert M, Sizoo B. Autism, adverse events, and trauma. In: Grabrucker AM, ed. Autism Spectrum Disorders. Exon Publications; 2021. Chapter 3. https://www.ncbi.nlm.nih.gov/books/NBK573608/ doi:
- 10.36255/exon publications. autism spectrum disorders. 2021. trauma
- 239. Quinton AMG, Ali D, Danese A, et al. The assessment and treatment of post-traumatic stress disorder in autistic people: A systematic review. Rev J Autism Dev Disord. 2024. https://doi.org/10.1007/s40489-024-00430-9
- 240. Kerns CM, Robins DL, Shattuck PT, Newschaffer CJ, Berkowitz SJ. Expert consensus regarding indicators of a traumatic reaction in autistic youth: a Delphi survey. J Child Psychol Psychiatry. 2023 Jan;64(1):50-58. doi: 10.1111/jcpp.13666. Epub 2022 Jul 11.
- 241. Rumball F, Happé F, Grey N. Experience of trauma and PTSD symptoms in autistic adults: Risk of PTSD development following DSM-5 and non-DSM-5 traumatic life events. Autism Res. 2020;13:12. p. 2122-2132. https://doi.org/10.1002/aur.2306

- 242. Telléus GK, Lauritsen MB, Rodrigo-Domingo M. Prevalence of various traumatic events including sexual trauma in a clinical sample of patients with an eating disorder. Front Psychol. 2021;8:12:687452. https://doi.org/10.3389/fpsyg.2021.687452
- 243. Gordon AR, Beccia AL, Egan N, Lipson SK. Intersecting gender identity and racial/ethnic inequities in eating disorder risk factors, symptoms, and diagnosis among U.S. college students: An intersectional multilevel analysis of individual heterogeneity and discriminatory accuracy. Int J Eat Disord. 2023. Advance online publication. doi: 10.1002/eat.24089
- 244. Urban B, Knutson D, Lazo-Salmeron W. Social-contextual influences on eating pathology in transgender and nonbinary adults of color. J LGBTQ Issues Couns. 2024;18. p. 143-159. doi: 10.1080/26924951.2024.2326614
- 245. Ward JH, Weir E, Allison C, Baron-Cohen S. Increased rates of chronic physical health conditions across all organ systems in autistic adolescents and adults. Mol Autism. 2023;20. doi: 10.1186/s13229-023-00565-2. PMID: 37730651; PMCID: PMC10510241.
- 246. Bradshaw P, Pickett C, van Driel ML, Brooker K, Urbanowicz A. Recognising, supporting and understanding autistic adults in general practice settings. Aust J Gen Pract. 2021;50:3. p. 126-130. https://doi.org/10.31128/ajgp-11-20-5722
- 247. Pan PY, Bölte S. The association between ADHD and physical health: a co-twin control study. Sci Rep. 2020 Dec 31;10:22388. https://doi.org/10.1038/s41598-020-78627-1
- 248. Csecs JLL, Iodice V, Rae CL, Brooke A, Simmons R, Quadt L, et al. Joint hypermobility links neurodivergence to dysautonomia and pain. Front Psychiatry. 2022 Feb 1;12:786916. https://doi.org/10.3389/fpsyt.2021.786916

- 249. Chen MH, Lan WH, Hsu JW, Huang KL, Su TP, Li CT, Lin WC, Tsai CF, Tsai SJ, Lee YC, Chen YS, Pan TL, Chang WH, Chen TJ, Bai YM. Risk of developing type 2 diabetes in adolescents and young adults with autism spectrum disorder: A nationwide longitudinal study. Diabetes Care. 2016 May;39:5. p. 788-793. https://doi.org/10.2337/dc15-1807
- 250. Hergüner S, Harmancı H, Toy H. Attention deficit-hyperactivity disorder symptoms in women with polycystic ovary syndrome. Int J Psychiatry Med. 2015;50:3. p. 317-325. https://doi.org/10.1177/0091217415610311
- 251. Gao M, Koupil I, Sjöqvist H, Karlsson H, Lalitkumar S, Dalman C, Kosidou K. Psychiatric comorbidity among women with endometriosis: Nationwide cohort study in Sweden. Am J Obstet Gynecol. 2020;223:3. https://doi.org/10.1016/j.ajog.2020.02.033
- 252. Xu G, Liu B, Yang W, Snetselaar LG, Chen M, Bao W, Strathearn L. Association of food allergy, respiratory allergy, and skin allergy with attention deficit/hyperactivity disorder among children. Nutrients. 2022;14:3:474. https://doi.org/10.3390/nu14030474
- 253. Chua RXY, Tay MJY, Ooi DSQ, Siah KTH, Tham EH, Shek LP, Meaney MJ, Broekman BFP, Loo EXL. Understanding the link between allergy and neurodevelopmental disorders: A current review of factors and mechanisms. Front Neurol. 2021;11:603571. https://doi.org/10.3389/fneur.2020.603571
- 254. Boonchooduang N, Louthrenoo O, Chattipakorn N, Chattipakorn SC. Possible links between gut-microbiota and attention-deficit/hyperactivity disorders in children and adolescents. Eur J Nutr. 2020;59:3391-3403.
- 255. Iglesias-Vázquez L, Van Ginkel Riba G, Arija V, Canals J. Composition of gut microbiota in children with autism spectrum disorder: A systematic review and meta-analysis. Nutrients. 2020;12:3.

- 256. Kim JY, Choi MJ, Ha S, Hwang J, Koyanagi A, Dragioti E, et al. Association between autism spectrum disorder and inflammatory bowel disease: A systematic review and meta-analysis. Autism Res. 2022;15:2. p. 340-352. https://doi.org/10.1002/aur.2656
- 257. Kamionkowski S, Shibli F, Ganocy S, Fass R. The relationship between gastroesophageal reflux disease and autism spectrum disorder in adult patients in the United States. Neurogastroenterol Motil. 2022;34:7. https://doi.org/10.1111/nmo.14295
- 258. Momen NC, Plana-Ripoll O, Bulik CM, McGrath JJ, Thornton LM, Yilmaz Z, Petersen LV. Comorbidity between types of eating disorder and general medical conditions. Br J Psychiatry. 2022;220:5. p. 279-286. doi:10.1192/bjp.2021.104
- 259. Carbone EA, D'Amato P, Vicchio G, De Fazio P, Segura-Garcia C. A systematic review on the role of microbiota in the pathogenesis and treatment of eating disorders. Eur Psychiatry. 2021;64:1.
- 260. Dhopatkar N, Keeler JL, Mutwalli H, Whelan K, Treasure J, Himmerich H. Gastrointestinal symptoms, gut microbiome, probiotics and prebiotics in anorexia nervosa: A review of mechanistic rationale and clinical evidence. Psychoneuroendocrinology. 2023;147:105959.
- 261. Di Lodovico L, Mondot S, Dore J, Mack I, Hanachi M, Gorwood P. Anorexia nervosa and gut microbiota: A systematic review and quantitative synthesis of pooled microbiological data. Prog Neuropsychopharmacol Biol Psychiatry. 2021;106:110114.
- 262. Hedman A, Breithaupt L, Hübel C, Thornton LM, Tillander A, Norring C, Birgegård A, Larsson H, Ludvigsson JF, Sävendahl L, Almqvist C, Bulik CM. Bidirectional relationship between eating disorders and autoimmune diseases. J Child Psychol Psychiatry. 2019;60:8. p. 803-812. https://doi.org/10.1111/jcpp.12958

- 263. Lalonde-Bester S, Malik M, Masoumi R, Ng K, Sidhu S, Ghosh M, Vine D. Prevalence and etiology of eating disorders in polycystic ovary syndrome: A scoping review. Adv Nutr. 2024;15:4:100193. doi:10.1016/j.advnut.2024.100193.
- 264. Elwyn R, Mitchell J, Kohn MR, Driver C, Hay P, Lagopoulos J, Hermens DF. Novel ketamine and zinc treatment for anorexia nervosa and the potential beneficial interactions with the gut microbiome. Neurosci Biobehav Rev. 2023;105122.
- 265. Murray SL, Holton KF. Post-traumatic stress disorder may set the neurobiological stage for eating disorders: A focus on glutamatergic dysfunction. Appetite. 2021;167:105599.
- 266. Rossi E, Cassioli E, Dani C, Marchesoni G, Monteleone AM, Wonderlich SA, et al. The maltreated eco-phenotype of eating disorders: a new diagnostic specifier? A systematic review of the evidence and comprehensive description. Neurosci Biobehav Rev. 2024;105619.
- 267. Fagundes CP, Glaser R, Kiecolt-Glaser JK. Stressful early life experiences and immune dysregulation across the lifespan. Brain Behav Immun. 2013 Jan;27(1):8-12. doi: 10.1016/j.bbi.2012.06.014. PMID: 22771426; PMCID: PMC3518756.
- 268. Yamanaka-Takaichi M, Mizukami Y, Sugawara K, Sunami K, Teranishi Y, Kira Y, Paus R, Tsuruta D. Stress and nasal allergy: corticotropin-releasing hormone stimulates mast cell degranulation and proliferation in human nasal mucosa. Int J Mol Sci. 2021;22(5):2773. https://doi.org/10.3390/ijms22052773.
- 269. Theoharides TC. The impact of psychological stress on mast cells. Ann Allergy Asthma Immunol. 2020;125:4. p. 388-392. https://doi.org/10.1016/j.anai.2020.07.007.
- 270. Kempuraj D, Selvakumar GP, Thangavel R, Ahmed ME, Zaheer S, Raikwar SP, Iyer SS, Bhagavan SM, Beladakere-Ramaswamy S, Zaheer A. Mast cell activation in brain injury,

stress, and post-traumatic stress disorder and Alzheimer's disease pathogenesis. Front Neurosci. 2017;11:703. https://doi.org/10.3389/fnins.2017.00703.

- 271. Cheng L, Liu J, Chen Z. The histaminergic system in neuropsychiatric disorders. Biomolecules. 2021;11:9:1345. https://doi.org/10.3390/biom11091345.
- 272. Weinstock LB, Nelson RM, Blitshteyn S. Neuropsychiatric manifestations of mast cell activation syndrome and response to mast-cell-directed treatment: a case series. J Pers Med. 2023;13:11:1562. https://doi.org/10.3390/jpm13111562.
- 273. Bennett FC, Molofsky AV. The immune system and psychiatric disease: a basic science perspective. Clin Exp Immunol. 2019;197:3. p. 294-307. https://doi.org/10.1111/cei.13334.
- 274. Soria V, Uribe J, Salvat-Pujol N, Palao D, Menchón JM, Labad J. Psychoneuroimmunology of mental disorders. Rev Psiquiatr Salud Ment (Engl Ed). 2017;11:2. p. 115-124. https://doi.org/10.1016/j.rpsmen.2017.07.002.
- 275. Zerwas S, Larsen JT, Petersen L, Thornton LM, Quaranta M, Koch SV, Pisetsky D, Mortensen PB, Bulik CM. Eating disorders, autoimmune, and autoinflammatory disease. Pediatrics. 2017;140:6.https://doi.org/10.1542/peds.2016-2089.
- 276. Sirufo MM, Magnanimi LM, Ginaldi L, De Martinis M. Anorexia nervosa and autoimmune comorbidities: A bidirectional route? CNS Neurosci Ther. 2022;28:12. p. 1921-1929. https://doi.org/10.1111/cns.13953.
- 277. Xu L, Lin W, Zheng Y, Chen J, Fang Z, Tan N, Hu W, Guo Y, Wang Y, Chen Z. An H2R-dependent medial septum histaminergic circuit mediates feeding behavior. Cell. 2022;32:9. p. 1937-1948. https://doi.org/10.1016/j.cub.2022.03.010.

- 278. Flik G, Folgering JH, Cremers TI, Westerink BH, Dremencov E. Interaction between brain histamine and serotonin, norepinephrine, and dopamine systems: in vivo microdialysis and electrophysiology study. J Mol Neurosci. 2015;56:2. p. 320-328. https://doi.org/10.1007/s12031-015-0536-3
- 279. Jin CY, Kalimo H, Panula P. The histaminergic system in human thalamus: correlation of innervation to receptor expression. Eur J Neurosci. 2002 Apr;15:7. p. 1125-1138. doi: 10.1046/j.1460-9568.2002.01951.x.
- 280. Sánchez-González MA, García-Cabezas MA, Rico B, Cavada C. The primate thalamus is a key target for brain dopamine. J Neurosci. 2005;25:26. p. 6076-6083. https://doi.org/10.1523/JNEUROSCI.0968-05.2005.
- 281. Wang KY, Tanimoto A, Yamada S, Guo X, Ding Y, Watanabe T, Kohno K, Hirano K, Tsukada H, Sasaguri Y. Histamine regulation in glucose and lipid metabolism via histamine receptors: model for nonalcoholic steatohepatitis in mice. Am J Pathol. 2010;177:2. p. 713-723. https://doi.org/10.2353/ajpath.2010.091198.
- 282. Khouma A, Moeini MM, Plamondon J, Richard D, Caron A, Michael NJ. Histaminergic regulation of food intake. Front Endocrinol (Lausanne). 2023 Jun 27;14:1202089. doi: 10.3389/fendo.2023.1202089.
- 283. Smolinska S, Winiarska E, Globinska A, Jutel M. Histamine: A mediator of intestinal disorders A review. Metabolites. 2022;12:10. https://doi.org/10.3390/metabo12100895.
- 285. Tabarean, IV. "Histamine receptor signaling in energy homeostasis." Neuropharmacology. 2016;13-9. doi:10.1016/j.neuropharm.2015.04.011

- 286. Mohammadi-Pilehdarboni H, Rasouli M. Histamine H1- and H2-receptors participate to provide metabolic energy differently. Fundam Clin Pharmacol. 2022;36:6. p. 1031-1037. https://doi.org/10.1111/fcp.12814
- 287. Keeler JL, Treasure J, Himmerich H. Immunological aspects of eating disorders. In: Robinson P, Wade T, Herpertz-Dahlmann B, Fernandez-Aranda F, Treasure J, Wonderlich S, eds. Eating Disorders. Springer; 2023. https://doi.org/10.1007/978-3-030-97416-9 45-1
- 288. Van Hoeken D, Hoek HW. Review of the burden of eating disorders: mortality, disability, costs, quality of life, and family burden. Curr Opin Psychiatry. 2020;33:6. p. 521-527.
- 289. Sala M, Keshishian A, Song S, Moskowitz R, Bulik CM, Roos CR, Levinson CA. Predictors of relapse in eating disorders: A meta-analysis. J Psychiatr Res. 2023;158. p. 281-299. https://doi.org/10.1016/j.jpsychires.2023.01.002
- 290. Wampold BE, Flückiger C, Del Re AC, Yulish NE, Frost ND, Pace BT, Goldberg SB, Miller SD, Baardseth TP, Laska KM, Hilsenroth MJ. In pursuit of truth: A critical examination of meta-analyses of cognitive behavior therapy. Psychother Res. 2017;27:1. p. 14-32. doi: 10.1080/10503307.2016.1249433.
- 291. Scheel AM, Schijen MRMJ, Lakens D. An excess of positive results: Comparing the standard psychology literature with registered reports. Adv Methods Pract Psychol Sci. 2021;4:2. https://doi.org/10.1177/25152459211007467
- 292. Haeffel GJ. Psychology needs to get tired of winning. R Soc Open Sci. 2022 Jun 22;9:6. https://doi.org/10.1098/rsos.220099

- 293. Bartoš F, Maier M, Shanks DR, Stanley TD, Sladekova M, Wagenmakers EJ. Meta-analyses in psychology often overestimate evidence for and size of effects. R Soc Open Sci. 2023;10:7. https://doi.org/10.1098/rsos.230224
- 294. Sandbank M, Bottema-Beutel K, Syu Y-C, Caldwell N, Feldman JI, Woynaroski T. Evidence-b(i)ased practice: Selective and inadequate reporting in early childhood autism intervention research. Autism. 2024. https://doi.org/10.1177/13623613241231624.
- 295. Shinohara K, Tajika A, Imai H, Takeshima N, Hayasaka Y, Furukawa TA. Protocol registration and selective outcome reporting in recent psychiatry trials: new antidepressants and cognitive behavioural therapies. Acta Psychiatr Scand. 2015;132:6. p. 489-498. https://doi.org/10.1111/acps.12502
- 296. Davis LC, Diianni AT, Drumheller SR, Elansary NN, D'Ambrozio GN, Herrawi F, Piper BJ, Cosgrove L. Undisclosed financial conflicts of interest in DSM-5-TR: cross sectional analysis. BMJ. 2024;384
- . https://doi.org/10.1136/bmj-2023-076902
- 297. Bottema-Beutel K, Crowley S. Pervasive undisclosed conflicts of interest in applied behavior analysis autism literature. Front Psychol. 2021 May 4;12:676303. https://doi.org/10.3389/fpsyg.2021.676303
- 298. Kekic M, Rose A, Baker C, Bayley D. Reduced self-harm on acute mental health wards following the implementation of a vision-based patient monitoring system: Evidence from five NHS trusts. J Psychiatr Ment Health Nurs. 2024. Advance online publication. https://doi.org/10.1111/jpm.13036

- 299. Klatte R, Strauss B, Flückiger C, Färber F, Rosendahl J. Defining and assessing adverse events and harmful effects in psychotherapy study protocols: A systematic review. Psychotherapy. 2023;60:1. p. 130-148. https://doi.org/10.1037/pst0000359
- 300. Curran J, Parry GD, Hardy GE, Darling J, Mason A-M, Chambers E. How does therapy harm? A model of adverse process using task analysis in the meta-synthesis of service users' experience. Front Psychol. 2019;10:347.

https://doi.org/10.3389/fpsyg.2019.00347

- 301. Dawson M, Fletcher-Watson S. When autism researchers disregard harms: A commentary. Autism. 2022;26:2. p. 564-566. https://doi.org/10.1177/13623613211031403
- 302. van Dis EAM, van Veen SC, Hagenaars MA, Batelaan NM, Bockting CLH, van den Heuvel RM, Cuijpers P, Engelhard IM. Long-term outcomes of cognitive behavioral therapy for anxiety-related disorders: A systematic review and meta-analysis. JAMA Psychiatry. 2020;77:3. p. 265-273. https://doi.org/10.1001/jamapsychiatry.2019.3986
- 303. Jubenville-Wood T, Nicholas DB, Weiss J, Cairns S. Facilitators and barriers in psychotherapy from the perspective of autistic adults: an enhanced critical incident study. Int J Qual Stud Health Well-being. 2023;18:1. https://doi.org/10.1080/17482631.2023.2278858
- 304. van Bree ESJ, Slof-Op't Landt MCT, van Furth EF. Predictors of recovery in eating disorders: A focus on different definitions. Int J Eat Disord. 2023;56:6. p. 1240-1245. https://doi.org/10.1002/eat.23950
- 305. Beygui NC, Cascio MA. Treating eating disorders in patients with autism: Should treatment standards be implemented on a spectrum? Pract Innov. 2022;7:4. p. 342-356. https://doi.org/10.1037/pri0000193

- 306. Nielsen S, Dobrescu SR, Dinkler L, Gillberg C, Gillberg C, Råstam M, et al. Effects of autism on 30-year outcome of anorexia nervosa. J Eat Disord. 2022;10:4. https://doi.org/10.1186/s40337-021-00518-1
- 307. Svedlund NE, Norring C, Ginsberg Y, von Hausswolff-Juhlin Y. Are treatment results for eating disorders affected by ADHD symptoms? A one-year follow-up of adult females. Eur Eat Disord Rev. 2018;26:4. p. 337-345. https://doi.org/10.1002/erv.2598
- 308. Svedlund NE, Norring C, Ginsberg Y, von Hausswolff-Juhlin Y. Symptoms of Attention Deficit Hyperactivity Disorder (ADHD) among adult eating disorder patients. BMC Psychiatry. 2017;17:19. https://doi.org/10.1186/s12888-016-1093-1
- 309. Babb C, Brede J, Jones CRG, Elliott M, Zanker C, Tchanturia K, Serpell L, Mandy W, Fox JRE. 'It's not that they don't want to access the support . . . it's the impact of the autism': The experience of eating disorder services from the perspective of autistic women, parents and healthcare professionals. Autism. 2021;25:5. p. 1409-1421.

https://doi.org/10.1177/1362361321991257

- 310. Kinnaird E, Tchanturia K. Looking beneath the surface: Distinguishing between common features in autism and anorexia nervosa. J Behav Cogn Ther. 2021;31:1. p. 3-13. https://doi.org/10.1016/j.jbct.2020.09.001
- 311. Junqueira DR, Phillips R, Zorzela L, Golder S, Loke Y, Moher D, Ioannidis JPA, Vohra S. Time to improve the reporting of harms in randomized controlled trials. J Clin Epidemiol. 2021;136. p. 216-220. https://doi.org/10.1016/j.jclinepi.2021.04.020
- 312. Parry GD, Crawford MJ, Duggan C. Iatrogenic harm from psychological therapies time to move on. Br J Psychiatry. 2016;208:3. p. 210-212. doi:10.1192/bjp.bp.115.163618

313. Jonsson U, Alaie I, Parling T, Arnberg FK. Reporting of harms in randomized controlled trials of psychological interventions for mental and behavioral disorders: a review of current practice. Contemp Clin Trials. 2014;38:1. p. 1-8. https://doi.org/10.1016/j.cct.2014.02.005

314. Papanikolaou PN, Churchill R, Wahlbeck K, Ioannidis JPA, The EU-PSI Project. Safety Reporting in Randomized Trials of Mental Health Interventions. Am J Psychiatry. 2004 Sep 1;161:9 p. 1692-1697. Available from: https://psychiatryonline.org/doi/full/10.1176/appi.ajp.161.9.1692s

- 315. Bottema-Beutel K, Crowley S, Sandbank M, Woynaroski TG. Adverse event reporting in intervention research for young autistic children. Autism. 2021;25:2. p. 322-335. doi:10.1177/1362361320965331
- 316. Monteleone AM, Pellegrino F, Croatto G, Carfagno M, Hilbert A, Treasure J, et al. Treatment of eating disorders: A systematic meta-review of meta-analyses and network meta-analyses. Neurosci Biobehav Rev. 2022;142:104857. doi:10.1016/j.neubiorev.2022.104857.
- 317. Linardon J, Wade TD, de la Piedad Garcia X, Brennan L. The efficacy of cognitive-behavioral therapy for eating disorders: A systematic review and meta-analysis. J Consult Clin Psychol. 2017;85:11. p. 1080-1094. doi:10.1037/ccp0000245.
- 318. Aves W. Escaping iatrogenic harm: A journey into mental health service avoidance. J Psychiatr Ment Health Nurs. 2024;00. p. 1-6. doi:10.1111/jpm.13020.
- 319. Brede J, Cage E, Trott J, Palmer L, Smith A, Serpell L, Mandy W, Russell A. "We Have to Try to Find a Way, a Clinical Bridge" autistic adults' experience of accessing and receiving support for mental health difficulties: A systematic review and thematic metasynthesis. Clin Psychol Rev. 2022;93:102131. doi:10.1016/j.cpr.2022.102131.

- 320. Camm-Crosbie L, Bradley L, Shaw R, Baron-Cohen S, Cassidy S. 'People like me don't get support': Autistic adults' experiences of support and treatment for mental health difficulties, self-injury and suicidality. Autism Int J Res Pract. 2019;23:6. p. 1431-1441. doi:10.1177/1362361318816053.
- 321. Lipinski S, Boegl K, Blanke ES, Suenkel U, Dziobek I. A blind spot in mental healthcare? Psychotherapists lack education and expertise for the support of adults on the autism spectrum. Autism Int J Res Pract. 2022;26:6. p. 1509-1521. doi:10.1177/13623613211057973.
- 322. Sara L. Livia Sara: Bridging the gap between autism and eating disorders. ED NEURODIVERSITY AU. Published December 20, 2023. Accessed July 11, 2024. Available from: https://www.edneuroaus.com/post/livia-sara-bridging-the-gap-between-autism-and-eating-disorders
- 323. Butterfly Foundation. Autism, eating disorders, and body image: Through the lens of sensory processing. Published April 1, 2021. Available from: https://butterfly.org.au/autism-eating-disorders-and-body-image-through-the-lens-of-sensory-processing/
- 324. Butterfly Foundation. Autism, eating disorders and me. Published January 12, 2022. Available from: https://butterfly.org.au/autism-eating-disorders-and-me/
- 325. Butterfly Foundation. Autism and eating disorders from the inside: My story. Butterfly Foundation. Published April 14, 2023. Available from: https://butterfly.org.au/autism-and-eating-disorders-from-the-inside-my-story/
- 326. Butterfly Foundation. Navigating eating disorder recovery as a neurodivergent individual. Published February 7, 2024. Available from: https://butterfly.org.au/navigating-eating-disorder-recovery-as-a-neurodivergent-individual/

- 327. Ritunnano R. Overcoming hermeneutical injustice in mental health: A role for critical phenomenology. J Br Soc Phenomenol. 2022;53:3. p. 243-260. doi:10.1080/00071773.2022.2031234.
- 328. Long Q, Jiang H. Qualitative research in health: value and visibility. Lancet Reg Health. 2023 May 10. Available from: https://doi.org/10.1016/j.lanwpc.2023.100790
- 329. Brause D. When therapy harms neurodivergent clients. Psychology Today. Published March 3, 2024. Available from: https://www.psychologytoday.com/au/blog/psychology-meets-neurodiversity/202402/when-therapy-harms-neurodivergent-clients
- 330. Downs J, Ayton A, Collins L, Baker S, Missen H, Ibrahim A. Untreatable or unable to treat? Creating more effective and accessible treatment for long-standing and severe eating disorders. Lancet Psychiatry. 2023;10:2. p. 146-154. doi:10.1016/S2215-0366(22)00400-X.
- 331. Elwyn R, Adams M, Sharpe SL, Silverstein S, LaMarre A, Downs J, et al. Discordant conceptualisations of eating disorder recovery and their influence on the construct of terminality. J Eat Disord. 2024;12:70. doi:10.1186/s40337-024-01016-w.
- 332. Curnow E, Rutherford M, Maciver D, Johnston L, Prior S, Boilson M, et al. Mental health in autistic adults: A rapid review of prevalence of psychiatric disorders and umbrella review of the effectiveness of interventions within a neurodiversity informed perspective. PLoS One. 2023;18:7. doi:10.1371/journal.pone.0288275.
- 333. Le Cunff A-L, Ellis Logan P, Ford R, Martis B-L, Mousset I, Sekibo J, Dommett E, Giampietro V. Co-Design for Participatory Neurodiversity Research: Collaborating With a Community Advisory Board to Design a Research Study. J Particip Res Methods. 2023;4:1. https://doi.org/10.35844/001c.66184.

- 334. Hu L. Towards a Critical Participatory Design Approach for the Neurodivergent: A Case Study for Students with ADHD. 2023.
- 335. den Houting J, Higgins J, Isaacs K, Mahony J, Pellicano E. From ivory tower to inclusion: Stakeholders' experiences of community engagement in Australian autism research. Front Psychol. 2022 Aug 24;13:876990.

https://doi.org/10.3389/fpsyg.2022.876990.

336. Gourdon-Kanhukamwe A, Kalandadze T, Yeung SK, Azevedo F, Iley B, Phan JM, Ramji AV, Shaw JJ, Zaneva M, Dokovova M, Hartmann H, Kapp SK, Warrington KL, Elsherif MM. Opening up understanding of neurodiversity: A call for applying participatory and open scholarship practices. Cogn Psychol Bull. 2023;1:8. p. 23-27. https://doi.org/10.53841/bpscog.2023.1.8.23

- 337. Cutler ES. Listening to Those with Lived Experience. In: Steingard S, editor. Critical Psychiatry. Springer; 2019. https://doi.org/10.1007/978-3-030-02732-2 8
- 338. Sonuga-Barke EJS, Chandler S, Lukito S, Kakoulidou M, Moore G, Cooper N, et al. Participatory translational science of neurodivergence: model for attention-deficit/hyperactivity disorder and autism research. Br J Psychiatry. 2024;224:4. p. 127-131. doi:10.1192/bjp.2023.151.
- 339. Pantazakos T, Vanaken GJ. Addressing the autism mental health crisis: the potential of phenomenology in neurodiversity-affirming clinical practices. Front Psychol. 2023;14:1225152. https://doi.org/10.3389/fpsyg.2023.1225152.
- 340. Keating CT. Participatory Autism Research: How Consultation Benefits Everyone. Front Psychol. 2021;12:713982. https://doi.org/10.3389/fpsyg.2021.713982.

- 341. Baker JH, Freestone D, Cai K, Silverstein S, Urban B, Steinberg D. Eating disorder clinical presentation and treatment outcomes by gender identity among children, adolescents, and young adults. J Adolesc Health. 2024. doi:10.1016/j.jadohealth.2023.11.015.
- 342. Wade TD, Shafran R, Cooper Z. Developing a protocol to address co-occurring mental health conditions in the treatment of eating disorders. Int J Eat Disord. 2024;57:6. p. 1291-1299. doi:10.1002/eat.24008.
- 343. Lai MC, Kassee C, Besney R, Bonato S, Hull L, Mandy W, Szatmari P, Ameis SH. Prevalence of co-occurring mental health diagnoses in the autism population: a systematic review and meta-analysis. Lancet Psychiatry. 2019;6:10. p. 819-829. https://doi.org/10.1016/S2215-0366(19)30289-5.
- 344. Hossain MM, Khan N, Sultana A, Ma P, McKyer ELJ, Ahmed HU, Purohit N. Prevalence of comorbid psychiatric disorders among people with autism spectrum disorder: An umbrella review of systematic reviews and meta-analyses. Psychiatry Res. 2020;287:112922. https://doi.org/10.1016/j.psychres.2020.112922.
- 345. Choi WS, Woo YS, Wang SM, Lim HK, Bahk WM. The prevalence of psychiatric comorbidities in adult ADHD compared with non-ADHD populations: A systematic literature review. PLoS One. 2022;17:11. https://doi.org/10.1371/journal.pone.0277175
- 346. Hughes-McCormack LA, Rydzewska E, Henderson A, MacIntyre C, Rintoul J, Cooper SA. Prevalence of mental health conditions and relationship with general health in a whole-country population of people with intellectual disabilities compared with the general population. BJPsych Open. 2017;3:5. p. 243-248. https://doi.org/10.1192/bjpo.bp.117.005462
- 347. Totsika V, Liew A, Absoud M, Adnams C, Emerson E. Mental health problems in children with intellectual disability. Lancet Child Adolesc Health. 2022;6:6. p. 432-444. https://doi.org/10.1016/S2352-4642(22)00067-0

- 348. Mitchison D, Broderstad AR, Burt A, Kvaløy K. Eating disorders in Indigenous peoples. In: Robinson P, Wade T, Herpertz-Dahlmann B, Fernandez-Aranda F, Treasure J, Wonderlich S, editors. Eating Disorders. Springer; 2023. p. 1-19. https://doi.org/10.1007/978-3-030-97416-9_19-1
- 349. Riddle MC, Safer JD. Medical considerations in the care of transgender and gender diverse patients with eating disorders. J Eat Disord. 2022;10:178. https://doi.org/10.1186/s40337-022-00699-3
- 350. Sackett DL, Rosenberg WM, Muir Gray JA, Haynes RB, Richardson WS. Evidence based medicine: what it is and what it isn't. BMJ. 1996;312:71. doi:10.1136/bmj.312.7023.7
- 351. Heruc G, Hurst K, Casey A, Fleming K, Freeman J, Fursland A, et al. ANZAED eating disorder treatment principles and general clinical practice and training standards. J Eat Disord. 2020;8:63. https://doi.org/10.1186/s40337-020-00341-0
- 352. Heruc G, Hart S, Stiles G, Fleming K, Casey A, Sutherland F, et al. ANZAED practice and training standards for dietitians providing eating disorder treatment. J Eat Disord. 2020;8:77. https://doi.org/10.1186/s40337-020-00334-z
- 353. Jeffrey S. Dietitian. Available from: https://eatingdisorderscarerhelpkit.com.au/wp-content/uploads/2019/10/RAVES-Model.pdf
- 354. Becker CB, Farrell NR, Waller G. Using exposure therapy for eating disorders. In: Smits JAJ, Jacquart J, Abramowitz J, Arch J, Margraf J, editors. Clinical Guide to Exposure Therapy: Beyond Phobias. Springer Nature Switzerland AG. 2022. pp. 277-297. https://doi.org/10.1007/978-3-031-04927-9_15

- 355. Dwyer P, Williams ZJ, Vukusic S, Saron CD, Rivera SM. Habituation of auditory responses in young autistic and neurotypical children. Autism Res. 2023;16:10. p. 1903-1923. https://doi.org/10.1002/aur.3022
- 356. Green SA, Hernandez L, Lawrence KE, Liu J, Tsang T, Yeargin J, Cummings K, Laugeson E, Dapretto M, Bookheimer SY. Distinct patterns of neural habituation and generalization in children and adolescents with autism with low and high sensory overresponsivity. Am J Psychiatry. 2019;176:12. p. 1010-1020. https://doi.org/10.1176/appi.ajp.2019.18121333
- 357. Walsh BT, Hagan KE, Lockwood C. A systematic review comparing atypical anorexia nervosa and anorexia nervosa. Int J Eat Disord. 2023;56:4. p. 798-820. https://doi.org/10.1002/eat.23856
- 358. Moskowitz L, Weiselberg E. Anorexia Nervosa/Atypical Anorexia Nervosa. Curr Probl Pediatr Adolesc Health Care. 2017;47:4. p. 70-84. doi:10.1016/j.cppeds.2017.02.003
- 359. Vo M, Golden N. Medical complications and management of atypical anorexia nervosa. J Eat Disord. 2022;10:196. https://doi.org/10.1186/s40337-022-00720-9
- 360. Ibrahim A, Ryan S, Viljoen D, et al. Integrated enhanced cognitive behavioural (I-CBTE) therapy significantly improves effectiveness of inpatient treatment of anorexia nervosa in real life settings. J Eat Disord. 2022;10:98. https://doi.org/10.1186/s40337-022-00620-y
- 361. Atwood ME, Friedman A. A systematic review of enhanced cognitive behavioral therapy (CBT-E) for eating disorders. Int J Eat Disord. 2020;53(3):311-330. https://doi.org/10.1002/eat.23206

- 362. Rankin R, Conti J, Ramjan L, Hay P. A systematic review of people's lived experiences of inpatient treatment for anorexia nervosa: Living in a "bubble." J Eat Disord. 2023;11:95. https://doi.org/10.1186/s40337-023-00820-0
- 363. Zugai JS, Gill K, Molloy L, Raeburn T, Ramjan L. The focus on weight in the inpatient care of anorexia nervosa: A qualitative investigation of consumer perspectives. Int J Ment Health Nurs. 2024;33:2. p. 388-396. https://doi.org/10.1111/inm.13242
- 364. Barko EB, Moorman SM. Weighing in: Qualitative explorations of weight restoration as recovery in anorexia nervosa. J Eat Disord. 2023;11:14. https://doi.org/10.1186/s40337-023-00736-9
- 365. Christian HP. "You're Just Looking at One Piece of the Puzzle... My Weight": A Phenomenological Examination of Diagnostic Crossover in Eating Disorders. Am J Qualitative Res. 2024;8:1. p. 57-70. https://doi.org/10.29333/ajqr/14084
- 366. Chapman R, Botha M. Neurodivergence-informed therapy. Dev Med Child Neurol. 2023;65:3. p. 310-317. https://doi.org/10.1111/dmcn.15384
- 367. Barbaro J. Neuroaffirming care values the strengths and differences of autistic people, those with ADHD or other profiles. Here's how. The Conversation. Published May 8, 2024. https://theconversation.com/neuroaffirming-care-values-the-strengths-and-differences-of-autistic-people-those-with-adhd-or-other-profiles-heres-how-227449
- 368. Hartman D, Day A, O'Donnell-Killen T, Doyle JK, Kavanagh M, Azevedo J. What does it mean to be neurodiversity affirmative? BPS. Published January 3, 2024. https://www.bps.org.uk/psychologist/what-does-it-mean-be-neurodiversity-affirmative

- 369. Mupotsa-Russell C. 'Complex' and 'diverse': Meaning-making and affirming practices as healing justice. Agenda. 2022;36:4. p. 189-197. https://doi.org/10.1080/10130950.2022.2149089
- 370. Doherty M, McCowan S, Shaw SCK. Autistic SPACE: a novel framework for meeting the needs of autistic people in healthcare settings. Br J Hosp Med. 2023;84:4. p. 1-9. doi:10.12968/hmed.2023.0006
- 371. McGreevy E, Quinn A, Law R, Botha M, Evans M, Rose K, Moyse R, Boyens T, Matejko M, Pavlopoulou G. An Experience Sensitive Approach to Care With and for Autistic Children and Young People in Clinical Services. J Humanist Psychol. 2024. doi:10.1177/00221678241232442
- 372. Patten KK, Murthi K, Onwumere DD, Skaletski EC, Little LM, Tomchek SD.

 Occupational Therapy Practice Guidelines for Autistic People Across the Lifespan. Am J

 Occup Ther. 2024 May 1;78:3:7803397010. doi: 10.5014/ajot.2024.078301.
- 373. Karlsson L, Råstam M, Wentz E. The SWedish Eating Assessment for Autism spectrum disorders (SWEAA)—Validation of a self-report questionnaire targeting eating disturbances within the autism spectrum. Res Dev Disabil. 2013;34:7. p. 2224-2233. doi:10.1016/j.ridd.2013.03.035
- 374. Gal E, Gal-Mishael R, Vissoker RE, Hedley D, Bury S, Stolar O. Eating Challenges in Children with Autism Spectrum Disorder: Development and Validation of the "Aut-Eat" Questionnaire (AEQ). J Autism Dev Disord. 2022;52. p. 811-822. https://doi.org/10.1007/s10803-021-04978-x
- 375. Lukens CT, Linscheid TR. Development and Validation of an Inventory to Assess Mealtime Behavior Problems in Children with Autism. J Autism Dev Disord. 2008;38. p. 342-352. https://doi.org/10.1007/s10803-007-0401-5

- 376. Adamson J, Brede J, Babb C, Serpell L, Jones CRG, Fox J, Mandy W. Towards identifying a method of screening for autism amongst women with restrictive eating disorders. Eur Eat Disord Rev. 2022 Sep;30:5. p. 592-603. doi: 10.1002/erv.2918.
- 377. PEACE. (n.d.). Peace pathway home. Available from: https://peacepathway.org/
- 378. Loomes R, Bryant-Waugh R. Widening the reach of family-based interventions for Anorexia Nervosa: autism-adaptations for children and adolescents. J Eat Disord. 2021;9:157. https://doi.org/10.1186/s40337-021-00511-8
- 379. Paynter J, Sommer K, Cook A. How can we make therapy better for Autistic Adults? Autistic Adults' Ratings of Helpfulness of Adaptations to Therapy. June 17, 2024. Available from: https://doi.org/10.31219/osf.io/7uaj6
- 380. Tchanturia K, Dandil Y, Li Z, Smith K, Leslie M, Byford S. A novel approach for autism spectrum condition patients with eating disorders: Analysis of treatment cost-savings. Eur Eat Disord Rev. 2021 May;29:3. p. 514-518. doi: 10.1002/erv.2760.
- 381. Dandil Y, Smith K, Adamson J, Tchanturia K. Individual cognitive remediation therapy benefits for patients with anorexia nervosa and high autistic features. Eur Eat Disord Rev. 2019 Nov;28(5):578-584. doi: 10.1002/erv.2707.
- 382. Loomes R, Bryant-Waugh R. Widening the reach of family-based interventions for anorexia nervosa: autism adaptations for children and adolescents. J Eat Disord. 2021;9:157. doi:10.1186/s40337-021-00511-8.
- 383. Millar C, Greenhill B. Adapted remote cognitive behavioural therapy for comfort eating with a woman with intellectual disabilities: Case report. J Eat Disord. 2022;10:29. doi:10.1186/s40337-022-00537-6.

- 384. Gulati S, Hameed B, Olusanya BO, Newton CR. Neurodiversity and humanism in autism: An LMIC health care setting perspective. Autism. 2023;27:5. p. 1173-1176. doi:10.1177/13623613231181477.
- 385. Wilkenfeld DA, McCarthy AM. Ethical concerns with applied behavior analysis for autism spectrum "disorder". Kennedy Inst Ethics J. 2020;30:1. p. 31-69. doi:10.1353/ken.2020.0000.
- 386. Anderson LK. Autistic experiences of applied behavior analysis. Autism. 2023;27:3. p. 737-750. doi:10.1177/13623613221118216.
- 387. Kupferstein H. Evidence of increased PTSD symptoms in autistics exposed to applied behavior analysis. Adv Autism. 2018;4:1. p. 19-29. doi:10.1108/aia-08-2017-0016 388. Sandoval-Norton, A. H., Shkedy, G., Shkedy, D., & Rushby, J. A. (2019). How much compliance is too much compliance: Is long-term ABA therapy abuse? Cogent Psychology,
- 389. Gorycki KA, Ruppel PR, Zane T, Navalta CP. Is long-term ABA therapy abusive: A response to Sandoval-Norton and Shkedy. Cogent Psychol. 2020;7:1. doi:10.1080/23311908.2020.1823615

6:1. https://doi.org/10.1080/23311908.2019.1641258

- 390. Hume R. Show me the real you: Enhanced expression of Rogerian conditions in therapeutic relationship building with autistic adults. Autism Adulthood. 2022;4:2. p. 151-163.
- 391. Catala A, Faucher L, Poirier P. Autism, epistemic injustice, and epistemic disablement: A relational account of epistemic agency. Synthese. 2021;199:3. p. 9013-9039.
- 392. Nicholson, C., Finlay, W. M. L., & Stagg, S. (2022). Self-determination and cooperation in supported mealtimes involving people with severe intellectual disabilities.

Disability and Rehabilitation, 45:17. p. 2741–2750.

https://doi.org/10.1080/09638288.2022.2104941

- 393. Redley M. Mealtime support for adults with intellectual disabilities: Understanding an everyday activity. J Appl Res Intellect Disabil. 2021;34:1. p. 111-117. doi:10.1111/jar.12790.
- 394. Davies J, Cooper K, Killick E, Sam E, Healy M, Thompson G, et al. Autistic identity: A systematic review of quantitative research. Autism Res. 2024;17:5. p. 874-897. doi:10.1002/aur.3105.
- 395. Cooper K, Russell AJ, Lei J, Smith LG. The impact of a positive autism identity and autistic community solidarity on social anxiety and mental health in autistic young people.

 Autism. 2023;27:3. p. 848-857. doi:10.1177/13623613221118351.
- 396. World Health Organization, United Nations. Mental health, human rights and legislation: Guidance and practice. World Health Organization. Published October 9, 2023. Available from: https://www.who.int/publications/i/item/9789240080737
- 397. Ma D, Su J, Wang H, Zhao Y, Li H, Li Y, et al. Sensory-based approaches in psychiatric care: A systematic mixed-methods review. J Adv Nurs. 2021;77:10. p. 3991-4004. doi:10.1111/jan.14884.
- 398. Wright L, Meredith P, Bennett S. Sensory approaches in psychiatric units: Patterns and influences of use in one Australian health region. Aust Occup Ther J. 2022;69:5. p. 559-573. doi:10.1111/1440-1630.12813.
- 399. Kandlur NR, Fernandes AC, Gerard SR, Rajiv S, Quadros S. Sensory modulation interventions for adults with mental illness: A scoping review. Hong Kong J Occup Ther. 2023;36:2. p. 57-68. doi:10.1177/15691861231204896.

- 400. Sutton D, Nicholson E. Sensory modulation in acute mental health wards: A qualitative study of staff and service user perspectives. Auckland, New Zealand: Te Pou o Te Whakaaro Nui; 2011.
- 401. Doroud N, Cappy M, Grant K, Scopelliti M, McKinstry C, McMahon D. Sensory rooms within mental health settings: A systematic scoping review. Occup Ther Ment Health. 2024. p. 1-21. doi:10.1080/0164212X.2024.2308290.
- 402. Greenwood E, Cooklin A, Barbaro J, Miller C. Autistic patients' experiences of the hospital setting: A scoping review. J Adv Nurs. 2024;80:3. p. 908-923. doi: 10.1111/jan.15880. Epub 2023 Sep 24.
- 403. Williams G, Corbyn J, Hart A. Improving the sensory environments of mental health in-patient facilities for autistic children and young people. Child Care Pract. 2023;29:1. p. 35-53. doi:10.1080/13575279.2022.2126437.